










INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : C07D 401/12, 405/14, A61K 31/44, A61P 7/02	A1	(11) International Publication Number: WO 00/31068 (43) International Publication Date: 2 June 2000 (02.06.00)
(21) International Application Number: PCT/IB99/02067 (22) International Filing Date: 17 November 1999 (17.11.99) (30) Priority Data: 09/196,921 19 November 1998 (19.11.98) US (71) Applicant: BERLEX LABORATORIES, INC. [US/US]; 15049 San Pablo Avenue, P.O. Box 4099, Richmond, CA 94804-0099 (US). (72) Inventor: PHILLIPS, Gary, B.; 3043 Shetland Drive, Pleasant Hill, CA 94523 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: POLYHYDROXYLATED HETEROCYCLIC DERIVATIVES AS ANTI-COAGULANTS		
(57) Abstract		
<p>This invention is directed to poly-hydroxylated monocyclic N-heterocyclic derivatives selected from formulae (I, II, III, IV, V, VI, VII), wherein Z¹, Z², R¹, R², R³, R⁴, R⁵, R⁶ and R⁷, are defined herein. These compounds are useful as anti-coagulants.</p>		
<div style="display: flex; flex-wrap: wrap; justify-content: space-around;"> <div style="text-align: center;">  <p>(I)</p> </div> <div style="text-align: center;">  <p>(II)</p> </div> <div style="text-align: center;">  <p>(III)</p> </div> <div style="text-align: center;">  <p>(IV)</p> </div> <div style="text-align: center;">  <p>(V)</p> </div> <div style="text-align: center;">  <p>(VI)</p> </div> <div style="text-align: center;">  <p>(VII)</p> </div> </div>		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav	TM	Turkmenistan
BF	Burkina Faso	GR	Greece		Republic of Macedonia	TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's	NZ	New Zealand		
CM	Cameroon		Republic of Korea	PL	Poland		
CN	China	KR	Republic of Korea	PT	Portugal		
CU	Cuba	KZ	Kazakhstan	RO	Romania		
CZ	Czech Republic	LC	Saint Lucia	RU	Russian Federation		
DE	Germany	LI	Liechtenstein	SD	Sudan		
DK	Denmark	LK	Sri Lanka	SE	Sweden		
EE	Estonia	LR	Liberia	SG	Singapore		

POLYHYDROXYLATED HETEROCYCLIC DERIVATIVES AS ANTI-COAGULANTS

Field of the Invention

5 The present invention is directed to polyhydroxylated monocyclic *N*-heterocyclic derivatives and their pharmaceutically acceptable salts, which inhibit the enzyme, factor Xa, thereby being useful as anti-coagulants. It also relates to pharmaceutical compositions containing the derivatives or their pharmaceutically acceptable salts, and methods of their use.

10

BACKGROUND OF THE INVENTION

Factor Xa is a member of the trypsin-like serine protease class of enzymes. A one-to-one binding of factors Xa and Va with calcium ions and phospholipid forms the prothrombinase complex which converts prothrombin to thrombin. Thrombin, in turn, converts fibrinogen to fibrin which polymerizes to form insoluble fibrin.

15

In the coagulation cascade, the prothrombinase complex is the convergent point of the intrinsic (surface activated) and extrinsic (vessel injury-tissue factor) pathways (*Biochemistry* (1991), Vol. 30, p. 10363; and *Cell* (1988), Vol. 53, pp. 505-518).

20 The model of the coagulation cascade has been refined further with the discovery of the mode of action of tissue factor pathway inhibitor (TFPI) (*Seminars in Hematology* (1992), Vol. 29, pp. 159-161). TFPI is a circulating multi-domain serine protease inhibitor with three Kunitz-type domains which competes with factor Va for free factor Xa. Once formed, the binary complex of factor Xa and TFPI becomes a potent inhibitor of the factor VIIa and tissue factor complex.

25

Factor Xa can be activated by two distinct complexes, by tissue factor-VIIa complex on the "Xa burst" pathway and by the factor IXa-VIIIa complex (TENase) of the "sustained Xa" pathway in the coagulation cascade. After vessel injury, the "Xa burst" pathway is activated *via* tissue factor (TF). Up regulation of the coagulation cascade occurs *via* increased factor Xa production *via* the "sustained Xa" pathway. Down regulation of the coagulation cascade occurs with the

30

formation of the factor Xa-TFPI complex, which not only removes factor Xa but also inhibits further factor formation *via* the "Xa burst" pathway. Therefore, the coagulation cascade is naturally regulated by factor Xa.

- 5 The primary advantage of inhibiting factor Xa over thrombin in order to prevent coagulation is the focal role of factor Xa versus the multiple functions of thrombin. Thrombin not only catalyzes the conversion of fibrinogen to fibrin, factor VIII to VIIIA, factor V to Va, and factor XI to XIa, but also activates platelets, is a monocyte chemotactic factor, and mitogen for lymphocytes and smooth muscle
- 10 cells. Thrombin activates protein C, the *in vivo* anti-coagulant inactivator of factors Va and VIIIA, when bound to thrombomodulin. In circulation, thrombin is rapidly inactivated by antithrombin III (ATIII) and heparin cofactor II (HCII) in a reaction which is catalyzed by heparin or other proteoglycan-associated
- 15 glycosaminoglycans, whereas thrombin in tissues is inactivated by the protease, nexin. Thrombin carries out its multiple cellular activation functions through a unique "tethered ligand" thrombin receptor (*Cell* (1991), Vol. 64, p. 1057), which requires the same anionic binding site and active site used in fibrinogen binding and cleavage and by thrombomodulin binding and protein C activation. Thus, a diverse group of *in vivo* molecular targets compete to bind thrombin and the
- 20 subsequent proteolytic events will have very different physiological consequences depending upon which cell type and which receptor, modulator, substrate or inhibitor binds thrombin.

- Published data with the proteins antistasin and tick anti-coagulant peptide (TAP)
- 25 demonstrate that factor Xa inhibitors are efficacious anti-coagulants (*Thrombosis and Haemostasis* (1992), Vol. 67, pp. 371-376; and *Science* (1990), Vol. 248, pp. 593-596).

- The active site of factor Xa can be blocked by either a mechanism-based or a tight
- 30 binding inhibitor (a tight binding inhibitor differs from a mechanism-based inhibitor by the lack of a covalent link between the enzyme and the inhibitor). Two types of mechanism-based inhibitors are known, reversible and irreversible, which are distinguished by ease of hydrolysis of the enzyme-inhibitor link (*Thrombosis Res*

(1992), Vol. 67, pp. 221-231; and *Trends Pharmacol. Sci.* (1987), Vol. 8, pp. 303-307). A series of guanidino compounds are examples of tight-binding inhibitors (*Thrombosis Res.* (1980), Vol. 19, pp. 339-349).

Arylsulfonyl-arginine-piperidine-carboxylic acid derivatives have also been shown
5 to be tight-binding inhibitors of thrombin (*Biochem.* (1984), Vol. 23, pp. 85-90), as
well as a series of arylamidine-containing compounds, including
3-amidinophenylaryl derivatives (*Thrombosis Res.* (1983), Vol. 29, pp. 635-642)
and bis(amidino)benzyl cycloketones (*Thrombosis Res.* (1980), Vol. 17,
pp. 545-548). However, these compounds demonstrate poor selectivity for factor
10 Xa.

Related Disclosures

European Published Patent Application 0 540 051 (Nagahara *et al.*) describes
aromatic amidine derivatives. These derivatives are stated to be capable of
15 showing a strong anticoagulant effect through reversible inhibition of factor Xa.

The synthesis of \forall, \forall' -bis(amidinobenzylidene)cycloalkanones and
 \forall, \forall' -bis(amidino-benzyl)cycloalkanones is described in *Pharmazie* (1977), Vol. 32,
No. 3, pp. 141-145. These compounds are disclosed as being serine protease
20 inhibitors.

U.S. Patent No. 5,451,700 (Morrissey *et al.*) describes amidino compounds.
These compounds are stated to be useful as selective LTB₄ receptor antagonists.

25 U.S. Patent No. 5,612,363 (Mohan *et al.*) describes *N,N*-di(aryl) cyclic urea
derivatives. These compounds are stated to be factor Xa inhibitors, thereby being
useful as anticoagulants.

U.S. Patent No. 5,633,381 (Dallas *et al.*) describes (Z,Z), (Z,E) and (E,Z) isomers
30 of substituted *bis*(phenylmethylene)cycloketones. These compounds are
disclosed as being factor Xa inhibitors, thereby being useful as anticoagulants.

PCT Published Patent Application WO/96/28427 (Buckman *et al.*) describes benzamidine derivatives. These compounds are stated to be factor Xa inhibitors, thereby being useful as anticoagulants.

- 5 PCT Published Patent Application WO/97/21437 (Arnaiz *et al.*) describes naphthyl-substituted benzimidazole derivatives. These compounds are disclosed as being factor Xa inhibitors, thereby being useful as anticoagulants.








- 10 PCT Published Patent Application WO/97/29067 (Kochanny *et al.*) describes benzamidine derivatives that are substituted by amino acid and hydroxy acid derivatives. These compounds are stated to be factor Xa inhibitors, thereby being useful as anticoagulants.

- 15 The above references, published patent applications and U.S. patents are herein incorporated in full by reference.

SUMMARY OF THE INVENTION

- 20 This invention is directed to compounds or their pharmaceutically acceptable salts which inhibit human factor Xa and are therefore useful as pharmacological agents for the treatment of disease-states characterized by thrombotic activity.

- Accordingly, in one aspect, this invention provides compounds selected from the group consisting of the following formulae:



wherein:

- 5 A is $-C(R^8)=$ or $-N=$ (where R^8 is hydrogen, alkyl or halo);
 Z^1 is $-O-$, $-N(R^9)-$, $-CH_2O-$ or $-S(O)_n-$ (where n is 0 to 2);
 Z^2 is $-O-$, $-N(R^9)-$, $-OCH_2-$ or $-S(O)_n-$ (where n is 0 to 2);
 R^1 and R^4 are each independently hydrogen, halo, alkyl, nitro, $-OR^9$, $-C(O)OR^9$,
 $-C(O)N(R^9)R^{10}$, $-N(R^9)R^{10}$, $-N(R^9)C(O)R^9$, or $-N(H)S(O)_2R^{11}$;
- 10 R^2 is $-C(NH)NH_2$, $-C(NH)N(H)OR^9$, $-C(NH)N(H)C(O)OR^{11}$, $-C(NH)N(H)C(O)R^9$,
 $-C(NH)N(H)S(O)_2R^{11}$, or $-C(NH)N(H)C(O)N(H)R^9$;

R^3 is hydrogen, halo, alkyl, haloalkyl, nitro, ureido, guanidino, $-OR^9$, $-C(NH)NH_2$,
 $-C(NH)N(H)OR^9$, $-C(O)N(R^9)R^{10}$, $-R^{12}-C(O)N(R^9)R^{10}$,
 $-CH(OH)C(O)N(R^9)R^{10}$, $-N(R^9)R^{10}$, $-R^{12}-N(R^9)R^{10}$, $-C(O)OR^9$, $-R^{12}-C(O)OR^9$,
 $-N(R^9)C(O)R^9$, (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl),
 5 (1,2)-imidazolyl (optionally substituted by alkyl), or (1,2)-imidazolinyl
 (optionally substituted by alkyl);

R^5 and R^6 are independently hydrogen, halo, alkyl, haloalkyl, nitro, $-N(R^9)R^{10}$,
 $-C(O)OR^9$, $-C(O)N(R^9)R^{10}$, $-C(O)N(R^9)CH_2C(O)N(R^9)R^{10}$,
 $-N(R^9)C(O)N(R^9)R^{10}$, $-N(R^9)C(O)R^{10}$, $-N(R^9)S(O)_2R^{11}$, or -
 10 $N(R^9)C(O)N(R^9)-CH_2C(O)N(R^9)R^{10}$;

R^7 is $-X-CH_2-[C(R^{13})H]_p-C(R^{13})H_2$ or $-X-C([C(R^{13})H]_p-C(R^{13})H_2)_2H$ where:


p is 0 to 5;

X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$;


and each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -
 15 $[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-[CH_2]_n-OC(O)R^9$, $-[CH_2]_n-$
 $SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$, $-[CH_2]_n-$
 $N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or $-[CH_2]_n-$
 $N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

or R^7 is selected from the group consisting of the following formulae:

20



and



where

q is 0 to 4;

25

p is 0 to 5;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -
 $[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-(CH_2)_n-OC(O)R^9$,
 $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$,
 $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0 to 2);

and

each R^{14} is $-\text{X-[CH}_2\text{]}_r$ or $-\text{X-CH}_2\text{-[C(R}^{13}\text{)H]}_r$, where:

each r is independently 0 to 5, and

each X is $-\text{O}-$, $-\text{S(O)}_n-$ (where n is 0 to 2), or $-\text{N(R}^9\text{)}-$; and

each R^{13} is independently $-\text{[CH(OR}^9\text{)]}_m\text{-CH}_2\text{-OR}^9$ (where m is 1 to 4),

$-\text{[CH}_2\text{]}_n\text{-OR}^9$, $-\text{[CH}_2\text{]}_n\text{-SR}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)R}^{10}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)R}^9$,

$-\text{[CH}_2\text{]}_n\text{-SC(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-OC(O)OR}^{11}$,

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)OR}^{11}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)N(R}^9\text{)R}^{10}$, or

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0

to 2);

each R^9 and R^{10} is independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl);

each R^{11} is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl); and

each R^{12} is independently an alkylene or alkylidene chain;

as a single stereoisomer or a mixture thereof; or a pharmaceutically acceptable salt thereof.

In another aspect, this invention provides compositions useful in treating a human having a disease-state characterized by thrombotic activity, which composition comprises a therapeutically effective amount of a compound of the invention as

described above, or a pharmaceutically acceptable salt thereof, and a pharmaceutically acceptable excipient.

In another aspect, this invention provides a method of treating a human having a
5 disease-state characterized by thrombotic activity, which method comprises administering to a human in need thereof a therapeutically effective amount of a compound of the invention as described above.

In another aspect, this invention provides a method of treating a human having a
10 disease-state alleviated by the inhibition of factor Xa, which method comprises administering to a human in need thereof a therapeutically effective amount of a compound of the invention as described above.

In another aspect, this invention provides a method of inhibiting human factor Xa
15 *in vitro* by the administration of a compound of the invention.

DETAILED DESCRIPTION OF THE INVENTION

Definitions

As used in the specification and appended claims, unless specified to the contrary,
20 the following terms have the meaning indicated:

"Alkyl" refers to a straight or branched hydrocarbon chain radical consisting solely of carbon and hydrogen atoms, containing no unsaturation, having from one to six carbon atoms, and which is attached to the rest of the molecule by a single bond,
25 e.g., methyl, ethyl, *n*-propyl, 1-methylethyl (*iso*-propyl), *n*-butyl, *n*-pentyl, 1,1-dimethylethyl (*t*-butyl), and the like.

"Alkoxy" refers to a radical of the formula -OR_a where R_a is alkyl as defined above, e.g., methoxy, ethoxy, *n*-propoxy, 1-methylethoxy (*iso*-propoxy), *n*-butoxy,
30 *n*-pentoxy, 1,1-dimethylethoxy (*t*-butoxy), and the like.

"Alkoxycarbonyl" refers to a radical of the formula $-C(O)OR_a$ where R_a is alkyl as defined above, e.g., methoxycarbonyl, ethoxycarbonyl, *n*-propoxycarbonyl, 1-methylethoxycarbonyl (*iso*-propoxycarbonyl), *n*-butoxycarbonyl, *n*-pentoxycarbonyl, 1,1-dimethylethoxycarbonyl (*t*-butoxycarbonyl), and the like.

5

"Alkylene chain" refers to straight or branched chain divalent radical consisting solely of carbon and hydrogen atoms, containing no unsaturation and having from one to six carbon atoms, e.g., methylene, ethylene, propylene, *n*-butylene and the like.

10

"Alkylidene chain" refers to a straight or branched chain unsaturated divalent radical consisting solely of carbon and hydrogen atoms, having from one to six carbon atoms, wherein the unsaturation is present only as double bonds and wherein a double bond can exist between the first carbon of the chain and the rest of the molecule, e.g., ethylidene, propylidene, *n*-butylidene, and the like.

15

"Aryl" refers to a phenyl or naphthyl radical.

"Aralkyl" refers to a radical of the formula $-R_aR_b$ where R_a is an alkyl radical, as defined above, substituted by R_b , an aryl radical, as defined above, e.g., benzyl.

20

"Aralkoxy" refers to a radical of the formula $-OR_c$ where R_c is an aralkyl radical as defined above, e.g., benzyloxy, and the like.

"Amidino" refers to the radical $-C(NH)NH_2$.

25

"Aminocarbonyl" refers to the radical $-C(O)NH_2$.

"Dialkylamino" refers to a radical of the formula $-N(R_a)R_a$ where each R_a is independently an alkyl radical as defined above, e.g., dimethylamino, methylethylamino, diethylamino, dipropylamino, ethylpropylamino, and the like.

30

"Dialkylaminocarbonyl" refers to a radical of the formula $-C(O)N(R_a)R_a$ where each R_a is independently an alkyl radical as defined above, e.g., dimethylaminocarbonyl, methylethylaminocarbonyl, diethylaminocarbonyl, dipropylaminocarbonyl, ethylpropylaminocarbonyl, and the like.

5

"Halo" refers to bromo, chloro, iodo or fluoro.

"Haloalkyl" refers to an alkyl radical, as defined above, that is substituted by one or more halo radicals, as defined above, e.g., trifluoromethyl, difluoromethyl, trichloromethyl, 2-trifluoroethyl, 1-fluoromethyl-2-fluoroethyl, 3-bromo-2-fluoropropyl, 1-bromomethyl-2-bromoethyl, and the like.

10

"(1,2)-Imidazolyl" refers to an imidazolyl radical attached at either the 1- or 2-position.

15

"(1,2)-Imidazolyl" refers to a 4,5-dihydroimidazolyl radical attached at either the 1- or the 2-position.

"Monoalkylamino" refers to a radical of the formula $-NHR_a$ where R_a is an alkyl radical as defined above, e.g., methylamino, ethylamino, propylamino, and the like.

20

"Monoalkylaminocarbonyl" refers to a radical of the formula $-C(O)NHR_a$ where R_a is an alkyl radical as defined above, e.g., methylaminocarbonyl, ethylaminocarbonyl, propylaminocarbonyl, and the like.

25

"(1,2)-Tetrahydropyrimidinyl" refers to a tetrahydropyrimidinyl radical attached at either the 1- or 2-position.

"Optional" or "optionally" means that the subsequently described event of circumstances may or may not occur, and that the description includes instances where said event or circumstance occurs and instances in which it does not. For example, "optionally substituted aryl" means that the aryl radical may or may not

30

be substituted and that the description includes both substituted aryl radicals and aryl radicals having no substitution.

"Pharmaceutically acceptable salt" includes both acid and base addition salts.

5

"Pharmaceutically acceptable acid addition salt" refers to those salts which retain the biological effectiveness and properties of the free bases, which are not biologically or otherwise undesirable, and which are formed with inorganic acids such as hydrochloric acid, hydrobromic acid, sulfuric acid, nitric acid, phosphoric acid and the like, and organic acids such as acetic acid, trifluoroacetic acid, propionic acid, glycolic acid, pyruvic acid, oxalic acid, maleic acid, malonic acid, succinic acid, fumaric acid, tartaric acid, citric acid, benzoic acid, cinnamic acid, mandelic acid, methanesulfonic acid, ethanesulfonic acid, *p*-toluenesulfonic acid, salicylic acid, and the like.

15

"Pharmaceutically acceptable base addition salt" refers to those salts which retain the biological effectiveness and properties of the free acids, which are not biologically or otherwise undesirable. These salts are prepared from addition of an inorganic base or an organic base to the free acid. Salts derived from inorganic bases include, but are not limited to, the sodium, potassium, lithium, ammonium, calcium, magnesium, iron, zinc, copper, manganese, aluminum salts and the like. Preferred inorganic salts are the ammonium, sodium, potassium, calcium, and magnesium salts. Salts derived from organic bases include, but are not limited to, salts of primary, secondary, and tertiary amines, substituted amines including naturally occurring substituted amines, cyclic amines and basic ion exchange resins, such as isopropylamine, trimethylamine, diethylamine, triethylamine, tripropylamine, ethanolamine, 2-dimethylaminoethanol, 2-diethylaminoethanol, dicyclohexylamine, lysine, arginine, histidine, caffeine, procaine, hydrabamine, choline, betaine, ethylenediamine, glucosamine, methylglucamine, theobromine, purines, piperazine, piperidine, *N*-ethylpiperidine, polyamine resins and the like. Particularly preferred organic bases are isopropylamine, diethylamine, ethanolamine, trimethylamine, dicyclohexylamine, choline and caffeine.

25

30

"Therapeutically effective amount" refers to that amount of a compound of the invention which, when administered to a human in need thereof, is sufficient to effect treatment, as defined below, for disease-states characterized by thrombotic activity. The amount of a compound of the invention which constitutes a

5 "therapeutically effective amount" will vary depending on the compound, the disease-state and its severity, and the age of the human to be treated, but can be determined routinely by one of ordinary skill in the art having regard to his own knowledge and to this disclosure.

10 "Treating" or "treatment" as used herein covers the treatment of a disease-state in a human, which disease-state is characterized by thrombotic activity, and includes:

- (i) preventing the disease-state from occurring in a human, in particular, when such human is predisposed to the disease-state but has not yet been
- 15 diagnosed as having it;
- (ii) inhibiting the disease-state, *i.e.*, arresting its development; or
- (iii) relieving the disease-state, *i.e.*, causing regression of the disease-state.

20


The yield of each of the reactions described herein is expressed as a percentage of the theoretical yield.

The compounds of the invention, or their pharmaceutically acceptable salts, may

25 have asymmetric carbon atoms, oxidized sulfur atoms or quaternized nitrogen atoms in their structure. The compounds of the invention and their pharmaceutically acceptable salts may therefore exist as single stereoisomers, racemates, and as mixtures of enantiomers and diastereomers. The compounds may also exist as geometric isomers. All such single stereoisomers, racemates


30 and mixtures thereof, and geometric isomers are intended to be within the scope of this invention.

The nomenclature used herein is a modified form of the I.U.P.A.C. system wherein the compounds of the invention are named as polyhydroxylated derivatives of benzamidine. For example, a compound of the invention selected from formula (I):




5

wherein A is -N=, Z¹ and Z² are both -O-; R¹ and R⁴ are both hydrogen, R² is -C(NH)NH₂, R³ is -C(O)N(R⁹)R¹⁰ (where R⁹ and R¹⁰ are both methyl), R⁵ and R⁶ are both fluoro, R⁷ is selected from the formula:




- 10 where -R¹⁴- is in the 2-position of the tetrahydrofuranyl moiety and is -X-(CH₂)_p- where X is -O- and p is 1; and q is three and one R¹³ is in the 3-position of the tetrahydrofuranyl moiety and is dimethylamino and another R¹³ is in the 4-position of the tetrahydrofuranyl moiety and is hydroxy, and another R¹³ is in the 5-position of the tetrahydrofuranyl moiety and is -CH₂-OH; *i.e.*, the compound of the following
- 15 formula:



is named herein as 3-[[3,5-difluoro-6-[3-(dimethylaminocarbonyl)phenoxy]-4-[[3-
 5 dimethylamino-4-hydroxy-5-(hydroxymethyl)tetrahydrofuran-2-yl]methoxy]pyridin-2-
 yl]oxy]benzamidine.

For purposes of this invention, parenthesis are used to denote substituents of a
 main atom and brackets are used to denote repeating sections of the substituent.
 For example, $-X-CH_2-[C(R^{13})H]_p-C(R^{13})H_2$ where p is 2 refers to the substituent of
 10 the formula:



Utility and Administration

15 A. Utility

The compounds of the invention are inhibitors of the serine protease, factor Xa,
 and are therefore useful in disease-states characterized by thrombotic activity
 based on factor Xa's role in the coagulation cascade (see Background of the
 Invention above). A primary indication for the compounds is prophylaxis for long
 20 term risk following myocardial infarction. Additional indications are prophylaxis of
 deep vein thrombosis (DVT) following orthopedic surgery or prophylaxis of
 selected patients following a transient ischemic attack. The compounds of the
 invention may also be useful for indications in which coumarin is currently used,
 such as for DVT or other types of surgical intervention such as coronary artery
 25 bypass graft and percutaneous transluminal coronary angioplasty. The
 compounds are also useful for the treatment of thrombotic complications
 associated with acute promyelocytic leukemia, diabetes, multiple myelomas,
 disseminated intravascular coagulation associated with septic shock, purpura
 fulminans associated infection, adult respiratory distress syndrome, unstable

angina, and thrombotic complications associated with aortic valve or vascular prosthesis. The compounds are also useful for prophylaxis for thrombotic diseases, in particular in patients who have a high risk of developing such disease.

- 5 In addition, the compounds of the invention are useful as *in vitro* and *in vivo* diagnostic reagents for selectively inhibiting factor Xa without inhibiting other components of the coagulation cascade.

10

B. Testing

The primary bioassays used to demonstrate the inhibitory effect of the compounds of the invention on factor Xa are simple chromogenic assays involving only serine protease, the compound of the invention to be tested, substrate and buffer (see, 15 *e.g.*, *Thrombosis Res.* (1979), Vol. 16, pp. 245-254). For example, four tissue human serine proteases can be used in the primary bioassay, free factor Xa, prothrombinase, thrombin (IIa) and tissue plasminogen activator (tPA). The assay for tPA has been successfully used before to demonstrate undesired side effects in the inhibition of the fibrinolytic process (see, *e.g.*, *J. Med. Chem.* (1993), Vol. 36, 20 pp. 314-319).

Another bioassay useful in demonstrating the utility of the compounds of the invention in inhibiting factor Xa demonstrates the potency of the compounds against free factor Xa in citrated plasma. For example, the anticoagulant efficacy 25 of the compounds of the invention will be tested using either the prothrombin time (PT), or activated partial thromboplastin time (aPTT) while selectivity of the compounds is checked with the thrombin clotting time (TCT) assay. Correlation of the K_i in the primary enzyme assay with the K_i for free factor Xa in citrated plasma will screen against compounds which interact with or are inactivated by other 30 plasma components. Correlation of the K_i with the extension of the PT is a necessary *in vitro* demonstration that potency in the free factor Xa inhibition assay translates into potency in a clinical coagulation assay. In addition, extension of the

PT in citrated plasma can be used to measure duration of action in subsequent pharmacodynamic studies.

For further information on assays to demonstrate the activity of the compounds of the invention, see R. Lottenberg *et al.*, *Methods in Enzymology* (1981), Vol. 80, pp. 341-361, and H. Ohno *et al.*, *Thrombosis Research* (1980), Vol. 19, pp. 579-588.

C. General Administration

Administration of the compounds of the invention, or their pharmaceutically acceptable salts, in pure form or in an appropriate pharmaceutical composition, can be carried out *via* any of the accepted modes of administration or agents for serving similar utilities. Thus, administration can be, for example, orally, nasally, parenterally, topically, transdermally, or rectally, in the form of solid, semi-solid, lyophilized powder, or liquid dosage forms, such as for example, tablets, suppositories, pills, soft elastic and hard gelatin capsules, powders, solutions, suspensions, or aerosols, or the like, preferably in unit dosage forms suitable for simple administration of precise dosages. The compositions will include a conventional pharmaceutical carrier or excipient and a compound of the invention as the/an active agent, and, in addition, may include other medicinal agents, pharmaceutical agents, carriers, adjuvants, *etc.*

Generally, depending on the intended mode of administration, the pharmaceutically acceptable compositions will contain about 1% to about 99% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, and 99% to 1% by weight of a suitable pharmaceutical excipient. Preferably, the composition will be about 5% to 75% by weight of a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, with the rest being suitable pharmaceutical excipients.

The preferred route of administration is oral, using a convenient daily dosage regimen which can be adjusted according to the degree of severity of the disease-state to be treated. For such oral administration, a pharmaceutically

acceptable composition containing a compound(s) of the invention, or a pharmaceutically acceptable salt thereof, is formed by the incorporation of any of the normally employed excipients, such as, for example, pharmaceutical grades of mannitol, lactose, starch, pregelatinized starch, magnesium stearate, sodium
5 saccharine, talcum, cellulose ether derivatives, glucose, gelatin, sucrose, citrate, propyl gallate, and the like. Such compositions take the form of solutions, suspensions, tablets, pills, capsules, powders, sustained release formulations and the like.

10 Preferably such compositions will take the form of capsule, caplet or tablet and therefore will also contain a diluent such as lactose, sucrose, dicalcium phosphate, and the like; a disintegrant such as croscarmellose sodium or derivatives thereof; a lubricant such as magnesium stearate and the like; and a binder such as a starch, gum acacia, polyvinylpyrrolidone, gelatin, cellulose ether derivatives, and the like.

15 The compounds of the invention, or their pharmaceutically acceptable salts, may also be formulated into a suppository using, for example, about 0.5% to about 50% active ingredient disposed in a carrier that slowly dissolves within the body, e.g., polyoxyethylene glycols and polyethylene glycols (PEG), e.g., PEG 1000
20 (96%) and PEG 4000 (4%).

Liquid pharmaceutically administrable compositions can, for example, be prepared by dissolving, dispersing, *etc.*, a compound(s) of the invention (about 0.5% to about 20%), or a pharmaceutically acceptable salt thereof, and optional
25 pharmaceutical adjuvants in a carrier, such as, for example, water, saline, aqueous dextrose, glycerol, ethanol and the like, to thereby form a solution or suspension.

If desired, a pharmaceutical composition of the invention may also contain minor
30 amounts of auxiliary substances such as wetting or emulsifying agents, pH buffering agents, antioxidants, and the like, such as, for example, citric acid, sorbitan monolaurate, triethanolamine oleate, butylated hydroxytoluene, *etc.*

Actual methods of preparing such dosage forms are known, or will be apparent, to those skilled in this art; for example, see *Remington's Pharmaceutical Sciences*, 18th Ed., (Mack Publishing Company, Easton, Pennsylvania, 1990). The composition to be administered will, in any event, contain a therapeutically effective amount of a compound of the invention, or a pharmaceutically acceptable salt thereof, for treatment of a disease-state alleviated by the inhibition of factor Xa in accordance with the teachings of this invention.


The compounds of the invention, or their pharmaceutically acceptable salts, are administered in a therapeutically effective amount which will vary depending upon a variety of factors including the activity of the specific compound employed; the metabolic stability and length of action of the compound; the age, body weight, general health, sex, and diet of the patient; the mode and time of administration; the rate of excretion; the drug combination; the severity of the particular disease-states; and the host undergoing therapy. Generally, a therapeutically effective daily dose is from about 0.14 mg to about 14.3 mg/kg of body weight per day of a compound of the invention, or a pharmaceutically acceptable salt thereof; preferably, from about 0.7 mg to about 10 mg/kg of body weight per day; and most preferably, from about 1.4 mg to about 7.2 mg/kg of body weight per day. For example, for administration to a 70 kg person, the dosage range would be from about 10 mg to about 1.0 gram per day of a compound of the invention, or a pharmaceutically acceptable salt thereof, preferably from about 50 mg to about 700 mg per day, and most preferably from about 100 mg to about 500 mg per day.

25

Preferred Embodiments

Of the compounds of the invention as set forth above in the Summary of the Invention, a preferred group of compounds are those compounds selected from formula (I):

30



as single stereoisomers or mixtures thereof; or as pharmaceutically acceptable
 5 salts thereof.

Of this group of compounds, a preferred subgroup of compounds are those compounds wherein:

A is -N=;

10 Z¹ is -O-, -CH₂O- or -S(O)_n- (where n is 1);

Z² is -O-, -OCH₂- or -S(O)_n- (where n is 1);

R¹ and R⁴ are each independently hydrogen, halo, alkyl or -OR⁹;

R² is -C(NH)NH₂, -C(NH)N(H)S(O)₂R¹¹ or -C(NH)N(H)C(O)N(H)R⁹;

R³ is ureido, guanidino, -N(R⁹)R¹⁰, -N(R⁹)C(O)R⁹, (1,2)-tetrahydropyrimidinyl

15 (optionally substituted by alkyl), (1,2)-imidazolyl (optionally substituted by alkyl) or (1,2)-imidazoliny (optionally substituted by alkyl);

R⁵ and R⁶ are independently hydrogen, halo, alkyl or haloalkyl;

R⁷ is -X-CH₂-[C(R¹³)H]_p-C(R¹³)H₂ where:

p is 0 to 5;

20 X is -O-, -S(O)_n- (where n is 0 to 2), or -N(R⁹)-;

and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -
 [CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -[CH₂]_n-OC(O)R⁹, -[CH₂]_n-
 SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-
 N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or -[CH₂]_n-
 25 N(R⁹)C(O)N(R⁹)R¹⁰ (where each n is independently 0 to 2);

each R⁹ and R¹⁰ is independently hydrogen, alkyl, aryl (optionally substituted by
 halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino,
 nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or

dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl); and

- 5 each R¹¹ is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
- 10 dialkylaminocarbonyl).

Of this subgroup of compounds, a preferred class of compounds are those compounds wherein:

- 15 A is -N=;
- Z¹ is -O-;
- Z² is -O-;
- R¹ is hydrogen or -OR⁹;
- R² is -C(NH)NH₂;
- 20 R³ is (1,2)-tetrahydropyrimidinyl (optionally substituted by methyl), (1,2)-imidazolyl (optionally substituted by methyl) or (1,2)-imidazolinyl (optionally substituted by methyl);
- R⁴ is hydrogen;
- R⁵ and R⁶ are each halo;
- 25 R⁷ is -X-CH₂-[C(R¹³)H]_p-C(R¹³)H₂ where:
- p is 0 to 5;
- X is -O-;
- and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -[CH₂]_n-OC(O)R⁹, -[CH₂]_n-SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or -[CH₂]_n-N(R⁹)C(O)N(R⁹)R¹⁰ (where each n is independently 0 to 2);
- 30 each R⁹ and R¹⁰ is independently hydrogen or alkyl; and

each R^{11} is alkyl or aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

5 Of this class of compounds, a preferred subclass of compounds are those compounds wherein:

A is $-N=$;

Z^1 is $-O-$;

Z^2 is $-O-$;

10 R^1 is $-OR^9$;

R^2 is $-C(NH)NH_2$;

R^3 is (1,2)-imidazolyl (optionally substituted by methyl) or (1,2)-imidazoliny (optionally substituted by methyl);

R^4 is hydrogen;

15 R^5 and R^6 are both fluoro;

R^7 is $-X-CH_2-[C(R^{13})H]_p-C(R^{13})H_2$ where:

p is 0 to 5;

X is $-O-$;

and each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -

20 $[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is independently 0 to 2); and

each R^9 and R^{10} is independently hydrogen or alkyl.

Of this subclass of compounds, preferred compounds are those compounds wherein

25 R^7 is $-O-CH_2-C(OH)H-C(OH)H_2$.

Of the preferred group of compounds of formula (I) as described above, another preferred subgroup of compounds are those compounds wherein:

A is $-N=$;

Z^1 is $-O-$, $-CH_2O-$ or $-S(O)_n-$ (where n is 0 to 2);

30 Z^2 is $-O-$, $-OCH_2-$ or $-S(O)_n-$ (where n is 0 to 2);

R^1 and R^4 are each independently hydrogen, halo or $-OR^9$;

R^2 is $-C(NH)NH_2$, $-C(NH)N(H)S(O)_2R^{11}$ or $-C(NH)N(H)C(O)N(H)R^9$;

R^3 is ureido, guanidino, $-N(R^9)R^{10}$, $-N(R^9)C(O)R^9$, (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl), (1,2)-imidazolyl (optionally substituted by alkyl), or (1,2)-imidazoliny (optionally substituted by alkyl);

R^5 and R^6 are independently hydrogen, halo, alkyl or haloalkyl;

5 R^7 is selected from the group consisting of the following formulae:



where

q is 0 to 4;

p is 0 to 5;

10 each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-(CH_2)_n-OC(O)R^9$, $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$, $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or $-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

15 and

each R^{14} is $-X-[CH_2]_r$ or $-X-CH_2-[C(R^{13})H]_r$, where:

each r is independently 0 to 5, and

each X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

20 each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-[CH_2]_n-OC(O)R^9$, $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$, $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or $-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

25 each R^9 and R^{10} is independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro,

carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl);

each R^{11} is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

Of this subgroup of compounds, a preferred class of compounds are those compounds wherein:

A is $-N=$;

Z^1 is $-O-$;

Z^2 is $-O-$;

R^1 is hydrogen or $-OR^9$;

R^2 is $-C(NH)NH_2$;

R^3 is (1,2)-tetrahydropyrimidinyl (optionally substituted by methyl), (1,2)-imidazolyl (optionally substituted by methyl), or (1,2)-imidazolyl (optionally substituted by methyl);

R^4 is hydrogen;

R^5 and R^6 are each halo;

R^7 is selected from the group consisting of the following formulae:



where

q is 0 to 4;


p is 0 to 5;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-(CH_2)_n-OC(O)R^9$,

$-\text{[CH}_2\text{]}_n\text{-SC(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-OC(O)OR}^{11}$,
 $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)OR}^{11}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)N(R}^9\text{)R}^{10}$, or
 $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0 to 2);
 and


- 5 each R^{14} is $-\text{X-[CH}_2\text{]}_r$ or $-\text{X-CH}_2\text{-[C(R}^{13}\text{)H]}_r$, where:
 each r is independently 0 to 5, and
 each X is $-\text{O-}$, $-\text{S(O)}_n$ - (where n is 0 to 2), or $-\text{N(R}^9\text{)-}$; and
 each R^{13} is independently $-\text{[CH(OR}^9\text{)]}_m\text{-CH}_2\text{-OR}^9$ (where m is 1 to 4),
 $-\text{[CH}_2\text{]}_n\text{-OR}^9$, $-\text{[CH}_2\text{]}_n\text{-SR}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)R}^{10}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)R}^9$,
 10 $-\text{[CH}_2\text{]}_n\text{-SC(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-OC(O)OR}^{11}$,
 $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)OR}^{11}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)N(R}^9\text{)R}^{10}$, or
 $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0
 to 2);
 each R^9 and R^{10} is independently hydrogen or alkyl; and
 15 each R^{11} is independently alkyl or aryl (optionally substituted by halo, alkyl,
 hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro,
 carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
 dialkylaminocarbonyl).

- 20 Of this class of compounds, a preferred subclass of compounds are those
 compounds wherein:
 A is $-\text{N=}$;
 Z^1 is $-\text{O-}$;
 Z^2 is $-\text{O-}$;
 25 R^1 is $-\text{OR}^9$;
 R^2 is $-\text{C(NH)NH}_2$;
 R^3 is (1,2)-imidazolyl (optionally substituted by methyl), or (1,2)-imidazoliny
 (optionally substituted by methyl);
 R^4 is hydrogen;
 30 R^5 and R^6 are both fluoro;
 R^7 is selected from the group consisting of the following formulae:



- 5 where
 q is 0 to 4;
 p is 0 to 5;
 each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-$
 $[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is
 10 independently 0 to 2); and
 each R^{14} is $-X-[CH_2]_r$ or $-X-CH_2-[C(R^{13})H]_r$, where:
 each r is independently 0 to 5, and
 each X is independently $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$;
 and
 15 each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4),
 $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$; and
 each R^9 and R^{10} is independently hydrogen or alkyl.

Of this subclass of compounds, a preferred set of compounds are those
 20 compounds wherein R^7 is:



25 where

q is 0 to 4;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$,
 $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is independently 0 to 2);

and

5 each R^{14} is $-X-[CH_2]_r$ or $-X-CH_2-[C(R^{13})H]_r$, where:


each r is independently 0 to 5, and

each X is independently $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4),
 $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$; and

10 each R^9 and R^{10} is independently hydrogen or alkyl.

Of this set of compounds, a preferred subset of compounds are those compounds wherein R^7 is:



where

15 q is 0 to 4;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$,
 $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is independently 0 to 2); and

each R^{14} is $-X-[CH_2]_r$ where:


r is 0 to 5, and

20 X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

each R^9 and R^{10} is independently hydrogen or alkyl.

Of this subset of compounds, preferred compounds are those compounds wherein:

25



where

q is 0 to 4;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$
or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is independently 0 to 2); and


5 each R^{14} is $-X-[CH_2]_r$ where:

r is 0 or 1, and

X is -O-; and

each R^9 and R^{10} is independently hydrogen or alkyl.

10 Of these preferred compounds, even more preferred compounds are those compounds wherein R^7 is:



where R^{14} is -O-; q is 3 and one R^{13} is in the 4-position of the tetrahydrofuranyl ring and is hydroxy, the second R^{13} is in the 2-position of the tetrahydrofuranyl ring and
15 is 2,3-dihydroxyethyl and the third R^{13} is in the 5-position of the tetrahydrofuranyl ring and is ethoxy.

Of the preferred compounds of the invention, the most preferred compounds are the following:

20 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[2-(1,2-dihydroxyethyl)-4-hydroxy-5-ethoxytetrahydrofuran-3-yl]oxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine; and
3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2,3-dihydroxypropoxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine.

25

Preparation of Compounds of The Invention


As a matter of convenience, the following description of the preparation of the compounds of the invention is directed to the preparation of compounds of formula (I) where A is -N=, Z^1 and Z^2 are both -O-, and R^2 is -C(NH)NH₂. It is understood,

however, that similar synthetic processes may be used to prepare other compounds of formula (I), (II), (III), (IV), (V), (VI), and (VII). It is also understood that in the following description, combinations of substituents and/or variables (e.g., R^9 and R^{10}) on the depicted formulae are permissible only if such combinations result in stable compounds.

Preparation of Compounds of Formula (Ia)

Compounds of formula (Ia) are compounds of Formula (I) of the invention wherein A is -N=, Z^1 and Z^2 are both -O-, and R^2 is -C(NH)NH₂. They are prepared as shown in the following REACTION SCHEME wherein R^1 , R^2 , R^3 , R^4 , R^5 , R^6 and R^7 are as described above in the Summary of the Invention, except that prior to Step 1, compounds of formula (B) which contain additional reactive hydroxy or amino groups may be treated with the appropriate oxygen- or nitrogen-protecting groups according to methods known to those skilled in the art, such as those described in Greene, T.W, *Protective Groups in Organic Synthesis* (1981), John Wiley & Sons, New York. Such protected compounds will be deprotected during the reaction conditions of Step 4 to form the desired R^7 substituent. Alternatively, compounds of formula (B) may already be present in an oxygen- or nitrogen-protected form and as a result of the reaction conditions of Step 4 may become deprotected to form the desired R^7 substituent.

REACTION SCHEME



- Compounds of formula (A), (B), (D), and (F) are commercially available, for example, from Aldrich Chemical Co., Sigma Chemical Co., or ICN Biomedicals, Inc, or may be prepared according to methods known to those skilled in the art. Compounds of formula (B) are also commercially available in protected forms, for example, from Aldrich Chemical Co. or Sigma Chemical Co.

In general, the compounds of formula (Ia) are prepared by first treating a compound of formula (A) with a compound of formula (B) in an aprotic solvent, for example, DMSO, in the presence of a base, for example, cesium carbonate, at 0°C to 80°C, preferably between 25°C to 60°C, for about 20 to 40 hours. The
5 compound of formula (C) is then isolated from the reaction mixture by standard techniques, such as extraction, filtration and *in vacuo* removal of solvent.

The resulting compound of formula (C) in an aprotic solvent, for example, acetonitrile, is treated with an equimolar amount of a compound of formula (D) in
10 the presence of a base, for example, cesium carbonate, at temperatures between about 20°C and 120°C, preferably at ambient temperature, for a period of time sufficient to complete the desired reaction as monitored by thin layer chromatography (TLC). The compound of formula (E) is then isolated from the reaction mixture by standard isolation techniques, such as extraction, *in vacuo*
15 removal of solvent, and flash chromatography.

The compound of formula (E) in an aprotic solvent, for example, DMSO, is then treated with an equimolar amount of a compound of formula (F) in the presence of a base, for example, cesium carbonate, at temperatures between about 20°C and
20 120°C, preferably at about 55°C, for a period of time sufficient to complete the desired reaction, for example, for about 24 hours. The reaction mixture is cooled to ambient temperature and the compound of formula (G) is then isolated from the reaction mixture through standard isolated techniques, such as extraction, *in vacuo* removal of solvent, and flash chromatography.

25 The compound of formula (G) is dissolved in an anhydrous alkanol, preferably ethanol, and then anhydrous mineral acid, preferably HCl, is added to the solution over a period of time sufficient to incorporate the acid into the solution while maintaining the reaction temperatures at about -78°C. After incorporation is
30 complete, the reaction vessel is sealed and the reaction mixture is allowed to warm to ambient temperature and stirred between 12 and 24 hours, preferably for about 16 hours. The solvent is removed *in vacuo* and the resulting residue is

dissolved in fresh anhydrous alkanol, preferably ethanol, and then treated with anhydrous ammonia (gas) at temperatures from between ambient temperature and about 100°C from about 1 to about 48 hours, preferably at about 60°C and for about 2 hours. The compound of formula (Ia) is then isolated from the reaction mixture by standard isolation techniques, for example, *in vacuo* removal of solvent and purification by high performance liquid chromatography (HPLC). During this last step, compounds of formula (G) where R⁷ is in an oxygen- or nitrogen-protected form are deprotected to form compounds of formula (Ia) where R⁷ is as defined above in the Summary of the Invention.

Alternatively, instead of treating the resulting residue above with anhydrous ammonia (gas), the resulting residue may be treated with a compound of the formula NH₂OR⁹ to afford the corresponding compound of formula (Ia) wherein R² is -C(NH)N(H)OR⁹.

Compounds of formula (Ia) wherein R³ is -C(NH)NH₂ or -C(NH)N(H)OR⁹ are produced from the corresponding cyano compounds in a similar manner as that described above for compound of formula (G).

Compounds of formula (Ia) wherein R¹, R³, R⁴, R⁵, R⁶, R⁹, R¹⁰ or R¹¹ contains an alkoxycarbonyl group or a -C(O)OR⁹ group where R⁹ is alkyl or aralkyl may also be treated under standard transesterification conditions with an alcohol of the formula R⁹OH where R⁹ is aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl) to produce compounds of the invention wherein R¹, R³, R⁴, R⁵, R⁶, R⁹, R¹⁰ or R¹¹ contains an aryloxycarbonyl group or a -C(O)OR⁹ group where R⁹ is aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

Compounds of formula (Ia) wherein R¹, R³, R⁴, R⁵, R⁶, R⁹, R¹⁰, or R¹¹ contains an aminocarbonyl group, a monoalkylaminocarbonyl group, a dialkylaminocarbonyl

- group, a $-C(O)N(R^9)R^{10}$ group or a $-C(O)OR^9$ group (where each R^9 or R^{10} is independently alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl) or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl)) may also be hydrolyzed under acidic conditions to prepare corresponding compounds of the invention where R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} , or R^{11} contains a carboxy or a $-C(O)OH$ group.
- 10 Compounds of formula (Ia) wherein R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} or R^{11} contains a carboxy group, an alkoxycarbonyl group, or a $-C(O)OR^9$ group where R^9 is hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or optionally substituted aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl) may also be amidated under standard amidation conditions to form the corresponding
- 15 compounds of formula (Ia) where R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} , or R^{11} contains an aminocarbonyl group, a monoalkylaminocarbonyl group, a dialkylaminocarbonyl group or a $-C(O)N(R^9)R^{10}$ group where R^9 and R^{10} are independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or optionally substituted aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).
- 20
- 25
- 30 Compounds of formula (Ia) where R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} , or R^{11} contains a nitro group may also be reduced under standard conditions to produce the corresponding compounds of formula (Ia) where R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} , or R^{11} contains an amino group, which may also be treated with the appropriate

alkylating agents or acylating agents to afford the corresponding compounds of formula (Ia) where R^1 , R^3 , R^4 , R^5 , R^6 , R^9 , R^{10} , or R^{11} contains a monoalkylamino group, a dialkylamino group, a $-N(R^9)R^{10}$ group or a $-N(R^9)C(O)R^9$ where each R^9 and R^{10} is independently hydrogen, alkyl, or aralkyl (optionally substituted by
5 halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

Compounds of formula (Ia) may also be further treated with the appropriate acid
10 halide, preferably acid chloride, or with the appropriate acid anhydride or an equivalent, to yield compounds of the invention wherein R^2 is $-C(NH)N(H)C(O)R^9$ where R^9 is hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
15 dialkylaminocarbonyl), or optionally substituted aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

20 Alternatively, compounds of formula (Ia) may further be treated with carbamoyl chlorides or their equivalents to yield compounds of the invention where R^2 is $-C(NH)N(H)C(O)OR^9$ where R^9 is hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
25 dialkylaminocarbonyl), or optionally substituted aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

30 Alternatively, compounds of formula (Ia) may also be further treated with compounds of the formula $R^{11}-S(O)_2$ -imidazole (where R^{11} is defined above in the Summary of the Invention) in a polar solvent, such as methylene chloride, at

ambient temperature to afford compounds of the invention where R^2 is -
 $C(NH)N(H)S(O)_2R^{11}$ where R^{11} is defined above in the Summary of the Invention.

Alternatively, compounds of formula (Ia) may be further treated with an
5 appropriately $N-R^9$ -substituted phenylcarbamate in a polar solvent, preferably
methylene chloride, at ambient temperature, for about 6 to 24 hours, preferably for
about 12 hours, to afford compounds of the invention where R^2 is -
 $C(NH)N(H)C(O)N(H)R^9$ where R^9 is defined above in the Summary of the
Invention.

10

Compounds of formula (Ia) which contain an unoxidized sulfur atom may be
oxidized with the appropriate oxidizing agent to produce compounds containing
oxidized sulfur (*i.e.*, $-S(O)_n-$ where n is 1 or 2).

* * * * *

15 The following specific preparations and examples are provided as a guide to assist
in the practice of the invention, and are not intended as a limitation on the scope of
the invention.

PREPARATION 1

20

Compounds of formula (C)

A. To pentafluoropyridine (2.0 g, 12 mmol) dissolved in DMSO was added
2,2-dimethyl-1,3-dioxolane-4-methanol (solketal) (1.5 mL, 12 mmol) and
 Cs_2CO_3 (3.8 g, 12 mmol). The slurry was heated in an oil bath at 60°C for
1 day. The reaction was partitioned with water and ethyl acetate, washed
25 with water and brine, dried (Na_2SO_4) and the solvent was removed to give
2,3,5,6-tetrafluoro-4-[[[(2,2-dimethyl)dioxolan-4-yl]methoxy]pyridine (3.2 g).

B. In a similar manner, the following compound of formula (C) was prepared:
2,3,5,6-tetrafluoro-4-[[[5-[(2,2-dimethyl)dioxolan-4-yl]-3a,5,6,6a-
30 tetrahydrofuro[2,3-*d*]-1,3-dioxol-6-yl]oxy]pyridine.

C. In a similar manner, the following oxygen- or nitrogen-protected compounds of formula (C) are prepared:

2,3,5,6-tetrafluoro-4-[2-(acetyl)amino-4-(methylthio)but-1-oxy]pyridine;

2,3,5,6-tetrafluoro-4-[(2-(acetyl)amino-3-hydroxyprop-1-yl)thio]pyridine;

5 2,3,5,6-tetrafluoro-4-[[2',2',2,2-tetramethyl-3a',4',7',7a'-tetrahydrospiro[1,3-dioxolane-4,6'-[6H-1,3]dioxolo[4,5-c]pyran-7'-yl]]oxy]pyridine;

2,3,5,6-tetrafluoro-4-[(4-methoxy-2,2-dimethyl-6-(trityloxy)methyl-3a,6,7,7a-tetrahydro-4H-1,3-dioxolo[4,5-c]pyran-7-yl)oxy]pyridine;

10 2,3,5,6-tetrafluoro-4-[(2,2-dimethyl-5-(2,2-dimethyl-1,3-dioxolan-4-yl)-1,3-dioxolan-4-yl)methoxy]pyridine;

2,3,5,6-tetrafluoro-4-[bis(2,2-dimethyl-1,3-dioxolan-4-yl)methoxy]pyridine;

2,3,5,6-tetrafluoro-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5H-bis[1,3]dioxolo[4,5-b:4',5'-d]pyran-3a-yl)methoxy]pyridine;

15 2,3,5,6-tetrafluoro-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5H-bis[1,3]dioxolo[4,5-b:4',5'-d]pyran-5-yl)methoxy]pyridine;

2,3,5,6-tetrafluoro-4-[(6-methoxy-2,2-dimethyl-3a,4,6,6a-tetrahydrofuro[3,4-d]-1,3-dioxol-4-yl)methoxy]pyridine; and

2,3,5,6-tetrafluoro-4-[(5-(((methoxycarbonyl)oxy)methyl)-2,2-dimethyl-3a,5,6,6a-tetrahydrofuro[2,3-d]-1,3-dioxol-6-yl)oxy]pyridine.

20

D. In a manner similar to that described above, other compounds of formula (C) may be prepared.

PREPARATION 2

25

Compounds of formula (E)

A. To 2,3,5,6-tetrafluoro-4-[(2,2-dimethyldioxolan-4-yl)methoxy]pyridine (3.2 g, 11 mmol) in CH₃CN (20 mL) was added Cs₂CO₃ (3.8 g, 12 mmol) and 3-hydroxy-4-(benzyloxy)benzonitrile (2.5 g, 11 mmol). After stirring for 24 hours the reaction was partitioned with water and ethyl acetate. The layers were separated, washed with water and brine, dried (Na₂SO₄), and the solvent was removed *in vacuo*. The residue was triturated with hexane/ethyl acetate (1/1) and the resulting solid was collected by

30

filtration to give 3-[[3,5,6-trifluoro-4-[(2,2-dimethyldioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile (3.0 g).

B. In a similar manner, the following compound of formula (E) was prepared:

5 3-[[3,5,6-trifluoro-4-[[5-[(2,2-dimethyl)dioxolan-4-yl]-3a,5,6,6a-tetrahydrofuro[2,3-*d*]-1,3-dioxol-6-yl]oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile.

C. In a similar manner, the following oxygen- or nitrogen-protected compounds of formula (E) are prepared:

10 3-[[3,5,6-trifluoro-4-[2-(acetyl)amino-4-(methylthio)but-1-oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[(2-(acetyl)amino-3-hydroxyprop-1-yl)thio]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[[2',2',2,2-tetramethyl-3a',4',7',7a'-tetrahydrospiro[1,3-
 15 doxolane-4,6'-[6*H*-1,3]dioxolo[4,5-*c*]pyran-7'-yl]]oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[(4-methoxy-2,2-dimethyl-6-(trityloxy)methyl-3a,6,7,7a-tetrahydro-4*H*-1,3-dioxolo[4,5-*c*]pyran-7-yl)oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 20 3-[[3,5,6-trifluoro-4-[(2,2-dimethyl-5-(2,2-dimethyl-1,3-dioxolan-4-yl)-1,3-dioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[*bis*(2,2-dimethyl-1,3-dioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5*H*-
 25 *bis*[1,3]dioxolo[4,5-*b*:4',5'-*d*]pyran-3a-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 3-[[3,5,6-trifluoro-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5*H*-
bis[1,3]dioxolo[4,5-*b*:4',5'-*d*]pyran-5-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
 30 3-[[3,5,6-trifluoro-4-[(6-methoxy-2,2-dimethyl-3a,4,6,6a-tetrahydrofuro[3,4-*d*]-1,3-dioxol-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile; and

3-[[3,5,6-trifluoro-4-[(5-(((methoxycarbonyl)oxy)methyl)-2,2-dimethyl-3a,5,6,6a-tetrahydrofuro[2,3-*d*]-1,3-dioxol-6-yl)oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile.

- 5 D. In a manner similar to that described above, other compounds of formula (E) may be prepared.

PREPARATION 3

Compounds of formula (G)

- 10 A. To 3-[[3,5,6-trifluoro-4-[(2,2-dimethyldioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile (1.0 g, 2.1 mmol) in DMSO (6 mL) was added Cs₂CO₃ (0.7 g, 2.1 mmol) and 2-(3-hydroxyphenyl)-1-methylimidazoline (0.4 g, 2.2 mmol). After stirring in an oil bath at 55°C for 24 hours, the reaction was partitioned with water and ethyl acetate.
- 15 The layers were separated, washed with water and brine, dried (Na₂SO₄), and the solvent was removed *in vacuo*. The residue was chromatographed on silica with CH₂Cl₂/MeOH/NH₄OH (240/10/1) to give 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2,2-dimethyldioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile (0.9 g).
- 20
- B. In a similar manner, the following compound of formula (G) was prepared: 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[5-[(2,2-dimethyl)dioxolan-4-yl]-3a,5,6,6a-tetrahydrofuro[2,3-*d*]-1,3-dioxol-6-yl]oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile.
- 25
- C. In a similar manner, the following oxygen-protected or nitrogen-protected compounds of formula (G) are prepared:
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[2-(acetyl)amino-4-(methylthio)but-1-oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 30 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2-(acetyl)amino-3-hydroxyprop-1-yl)thio]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;

- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[2',2',2,2-tetramethyl-3a',4',7',7a'-tetrahydrospiro[1,3-dioxolane-4,6'-[6*H*-1,3]dioxolo[4,5-*c*]pyran-7'-yl]]oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 5 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(4-methoxy-2,2-dimethyl-6-(trityloxy)methyl-3a,6,7,7a-tetrahydro-4*H*-1,3-dioxolo[4,5-*c*]pyran-7-yl)oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2,2-dimethyl-5-(2,2-dimethyl-1,3-dioxolan-4-yl)-1,3-dioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 10 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-*bis*(2,2-dimethyl-1,3-dioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5*H*-*bis*[1,3]dioxolo[4,5-*b*:4',5'-*d*]pyran-3a-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 15 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2,2,7,7-tetramethyl-3a,5a,8a,8b-tetrahydro-5*H*-*bis*[1,3]dioxolo[4,5-*b*:4',5'-*d*]pyran-5-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(6-methoxy-2,2-dimethyl-3a,4,6,6a-tetrahydrofuro[3,4-*d*]-1,3-dioxol-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile; and
- 20 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(5-(((methoxycarbonyl)oxy)methyl)-2,2-dimethyl-3a,5,6,6a-tetrahydrofuro[2,3-*d*]-1,3-dioxol-6-yl)oxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile.
- 25 D. In a manner similar to that described above, other compounds of formula (G) may be prepared.

EXAMPLE 1

Compounds of formula (I)

- A. To 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2,2-dimethyldioxolan-4-yl)methoxy]pyridin-2-yl]oxy]-4-(benzyloxy)benzonitrile
5 (0.9 g, 1.4 mmol), dissolved in ethanol (6 mL) and cooled in a dry ice/isopropanol bath, was bubbled HCl (g). After the solution was saturated, the reaction flask was sealed and the reaction mixture was allowed to warm to ambient temperature and was stirred for 18 hours. The solvent was removed *in vacuo* and the residue was triturated with
10 ether. The ether was decanted off and the residue was dissolved in ethanol (6 mL). The solution was cooled in a dry ice/isopropanol bath and ammonia (g) was bubbled into the solution. The reaction flask was sealed and heated in an oil bath at 60°C for 2 hours. The solvent was removed *in vacuo* and the residue was purified by HPLC on a C18 Dynamax
15 column with a 10-40% acetonitrile in water gradient with 0.1% trifluoroacetic acid to give 0.76 g of 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2,3-dihydroxypropoxy)pyridin-2-yl]oxy]-4-hydroxybenzamidinium as a pure trifluoroacetic acid salt; NMR (DMSO- d_6) 10.2 (s,2), 9.0 (s,2), 8.9 (s,2), 7.5 (m,3), 7.4 (m,3), 7.0 (d,1), 4.55
20 (m,1), 4.4 (m,1), 4.0 (m,2), 3.9 (m,2), 3.8 (m,1), 3.45 (d,2), 2.95 (s,3) ppm.
- B. In a similar manner, the following compound of formula (I) was made:
3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[2-(1,2-dihydroxyethyl)-
4-hydroxy-5-ethoxytetrahydrofuran-3-yl]oxy]pyridin-2-yl]oxy]-4-
25 hydroxybenzamidinium, trifluoroacetic acid salt; NMR (DMSO- d_6) 11.1 (s,1), 10.3 (s,1), 9.0 (s,2), 8.9 (s,2), 7.5 (m,3), 7.3 (m,3), 7.0 (d,1), 5.5 (m,1), 5.3 (m,1), 4.75 (m,2), 4.0 (m,2), 3.85 (m,2), 3.6 (m,4), 3.5 (m,3), 2.92 (s,3), 1.12 (t,3) ppm.
- C. In a similar manner, the following compounds of the invention are
30 prepared:

- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2-amino-4-(methylthio)but-1-oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 5 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((2-amino-3-hydroxyprop-1-yl)thio)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((4,5-dihydroxy-2-ethoxy-2-(hydroxymethyl)tetrahydropyran-3-yl)oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 10 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((4,5-dihydroxy-2-hydroxymethyl-6-ethoxytetrahydropyran-3-yl)oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((2,3,4,5-tetrahydroxypent-1-yl)oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 15 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((1,2,4,5-tetrahydroxypent-3-yl)oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(2-ethoxy-3,4,5-trihydroxytetrahydropyran-2-yl)methoxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 20 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(6-ethoxy-3,4,5-trihydroxytetrahydropyran-2-yl)methoxy]pyridine-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt;
- 25 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(3,4-dihydroxy-5-ethoxytetrahydrofuran-2-yl)methoxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt; and
- 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[(5-ethoxy-4-hydroxy-2-hydroxymethyltetrahydrofuran-3-yl)oxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine, trifluoroacetic acid salt.
- 30

D. In a manner similar to that described above, other compounds of the invention may be prepared.

EXAMPLE 2

This example illustrates the preparation of representative pharmaceutical compositions for oral administration containing a compound of the invention, or a pharmaceutically acceptable salt thereof, e.g., 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[2-(1,2-dihydroxyethyl)-4-hydroxy-5-ethoxytetrahydrofuran-3-yl]oxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine:

	A. <u>Ingredients</u>	<u>% wt./wt.</u>
	Compound of the invention	20.0%
10	Lactose	79.5%
	Magnesium stearate	0.5%

The above ingredients are mixed and dispensed into hard-shell gelatin capsules containing 100 mg each, one capsule would approximate a total daily dosage.

	B. <u>Ingredients</u>	<u>% wt./wt.</u>
15	Compound of the invention	20.0%
	Magnesium stearate	0.9%
	Starch	8.6%
	Lactose	69.6%
	PVP (polyvinylpyrrolidone)	0.9%

20 The above ingredients with the exception of the magnesium stearate are combined and granulated using water as a granulating liquid. The formulation is then dried, mixed with the magnesium stearate and formed into tablets with an appropriate tableting machine.

C. Ingredients

Compound of the invention	0.1 g
Propylene glycol	20.0 g
Polyethylene glycol 400	20.0 g
Polysorbate 80	1.0 g
Water	q.s. 100 mL

The compound of the invention is dissolved in propylene glycol, polyethylene glycol 400 and polysorbate 80. A sufficient quantity of water is then added with stirring to provide 100 mL of the solution which is filtered and bottled.

D. Ingredients% wt./wt.

Compound of the invention	20.0%
Peanut Oil	78.0%
Span 60	2.0%

The above ingredients are melted, mixed and filled into soft elastic capsules.

E. Ingredients% wt./wt.

Compound of the invention	1.0%
Methyl or carboxymethyl cellulose	2.0%
0.9% saline	q.s. 100 mL

The compound of the invention is dissolved in the cellulose/saline solution, filtered and bottled for use.

EXAMPLE 3

This example illustrates the preparation of a representative pharmaceutical formulation for parenteral administration containing a compound of the invention, or a pharmaceutically acceptable salt thereof, e.g., 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2,3-dihydroxypropoxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine:

Ingredients

Compound of the invention	0.02 g
Propylene glycol	20.0 g
Polyethylene glycol 400	20.0 g
Polysorbate 80	1.0 g
0.9% Saline solution	q.s. 100 mL

The compound of the invention is dissolved in propylene glycol, polyethylene glycol 400 and polysorbate 80. A sufficient quantity of 0.9% saline solution is then added with stirring to provide 100 mL of the I.V. solution which is filtered
5 through a 0.2 m membrane filter and packaged under sterile conditions.

EXAMPLE 4

This example illustrates the preparation of a representative pharmaceutical composition in suppository form containing a compound of the invention, or a
10 pharmaceutically acceptable salt thereof, e.g., 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2,3-dihydroxypropoxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine:

	<u>Ingredients</u>	<u>% wt./wt.</u>
	Compound of the invention	1.0%
15	Polyethylene glycol 1000	74.5%
	Polyethylene glycol 4000	24.5%

The ingredients are melted together and mixed on a steam bath, and poured into molds containing 2.5 g total weight.

20 EXAMPLE 5

This example illustrates the preparation of a representative pharmaceutical formulation for insufflation containing a compound of the invention, or a
pharmaceutically acceptable salt thereof, e.g., 3-[[3,4-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2-amino-4-(methylthio)but-1-oxy)pyridin-2-
25 yl]oxy]-4-hydroxybenzamidine:

<u>Ingredients</u>	<u>% wt./wt.</u>
Micronized compound of the invention	1.0%
Micronized lactose	99.0%

The ingredients are milled, mixed, and packaged in an insufflator equipped with
30 a dosing pump.

EXAMPLE 6

This example illustrates the preparation of a representative pharmaceutical formulation in nebulized form containing a compound of the invention, or a pharmaceutically acceptable salt thereof, e.g., 3-[[3,4-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2-amino-4-(methylthio)but-1-oxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine:

	<u>Ingredients</u>	<u>% wt./wt.</u>
	Compound of the invention	0.005%
	Water	89.995%
10	Ethanol	10.000%

The compound of the invention is dissolved in ethanol and blended with water. The formulation is then packaged in a nebulizer equipped with a dosing pump.

EXAMPLE 7

15 This example illustrates the preparation of a representative pharmaceutical formulation in aerosol form containing a compound of the invention, or a pharmaceutically acceptable salt thereof, e.g., 3-[[3,4-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-((2-amino-3-hydroxyprop-1-yl)thio)pyridin-2-yl]oxy]-4-hydroxybenzamidine:

	<u>Ingredients</u>	<u>% wt./wt.</u>
20	Compound of the invention	0.10%
	Propellant 11/12	98.90%
	Oleic acid	1.00%

The compound of the invention is dispersed in oleic acid and the propellants.

25 The resulting mixture is then poured into an aerosol container fitted with a metering valve.

EXAMPLE 8

(*In vitro* assay for Factor Xa and Thrombin)

30 This assay demonstrates the activity of the compounds of the invention towards factor Xa, thrombin and tissue plasminogen activator. The activities were determined as an initial rate of cleavage of the peptide *p*-nitroanilide by the

enzyme. The cleavage product, *p*-nitroaniline, absorbs at 405 nm with a molar extinction coefficient of $9920 \text{ M}^{-1}\text{cm}^{-1}$.

Reagents and Solutions:

Dimethyl sulfoxide (DMSO) (Baker analyzed grade).

5 Assay buffer:

50 mM TrisHCl, 150 mM NaCl, 2.5 mM CaCl_2 , and
0.1% polyethylene glycol 6000, pH 7.5.

Enzymes (Enzyme Research Lab.):

1. Human factor Xa stock solution: 0.281 mg/mL in assay buffer, stored at
10 -80°C (working solution (2X): 106 ng/mL or 2 nM in assay buffer, prepare
prior to use).
2. Human thrombin stock solution: Concentration as specified by the
supplier, stored at -80°C (working solution (2X): 1200 ng/mL or 32 nM in
assay buffer, prepare prior to use).
- 15 3. Human tissue plasminogen activator (tPA) (Two chains, Sigma or
American Diagnostica Inc.) stock solution: Concentration as specified by
the supplier, stored at -80°C (working solution (2X): 1361 ng/mL or 20 nM
in assay buffer, prepare prior to use).

Chromogenic substrates (Pharmacia Hepar Inc.):

- 20 1. S2222 (FXa assay) stock solution: 6 mM in deionized H_2O , store at 4°C
(working solution (4X): $656 \mu\text{M}$ in assay buffer).
2. S2302 (Thrombin assay) stock solution: 10 mM in deionized H_2O , stored
at 4°C (working solution (4X): $1200 \mu\text{M}$ in assay buffer).
3. S2288 (tPA assay) stock solution: 10 mM in deionized H_2O , stored at 4°C
25 (working solution (4X): $1484 \mu\text{M}$ in assay buffer for Sigma tPA, or 1120
 μM for American Diagnostica tPA).

Standard inhibitor compound stock solution:

5 mM in DMSO, stored at -20°C .

Test compounds (compounds of the invention) stock solutions:

- 30 10 mM in DMSO, stored at -20°C .

Assay procedure:

Assays were performed in 96-well microtiter plates in a total volume of 200 μ L. Assay components were in final concentration of 50 mM TrisHCl, 150 mM NaCl, 2.5 mM CaCl_2 , 0.1% polyethylene glycol 6000, pH 7.5, in the absence or
5 presence of the standard inhibitor or the test compounds and enzyme and substrate at following concentrations: (1) 1 nM factor Xa (0.1 nM or 0.2 nM factor Xa for compounds with K_i Xa in low picomolar range) and 164 μ M S2222; (2) 16 nM thrombin and 300 μ M S2302; and (3) 10 nM tPA and 371 μ M or 280 μ M S2288. Concentrations of the standard inhibitor compound in the assay were
10 from 5 μ M to 0.021 μ M in 1 to 3 dilution. Concentration of the test compounds in the assay typically were from 10 μ M to 0.041 μ M in 1 to 3 dilution. For potent test compounds, the concentrations used in the factor Xa assay were further diluted 100 fold (100 nM to 0.41 nM) or 1000 fold (10 nM to 0.041 nM). All substrate concentrations used are equal to their K_m values under the present
15 assay conditions. Assays were performed at ambient temperature.

The first step in the assay was the preparation of 10 mM test compound stock solutions in DMSO (for potent test compounds, 10 mM stock solutions were further diluted to 0.1 or 0.01 mM for the factor Xa assay), followed by the
20 preparation of test compound working solutions (4X) by a serial dilutions of 10 mM stock solutions with Biomek 1000 in 96 deep well plates as follows:
(a) Prepare a 40 μ M working solution by diluting the 10 mM stock 1 to 250 in assay buffer in 2 steps: 1 to 100, and 1 to 2.5.
25 (b) Make another five serial dilutions (1:3) of the 40 μ M solution (600 μ L for each concentration). A total of six diluted test compound solutions were used in the assay.

Standard inhibitor compound (5 mM stock) or DMSO (control) went through the
30 same dilution steps as those described above for test compounds.

The next step in the assay was to dispense 50 μ L of the test compound working

solutions (4X) (from 40 μ M to 0.164 μ M) in duplicate to microtiter plates with Biomek. To this was added 100 μ L of enzyme working solution (2X) with Biomek. The resulting solutions were incubated at ambient temperature for 10 minutes.

.5

To the solutions was added 50 μ L of substrate working solution (4X) with Biomek.

The enzyme kinetics were measured at 405 nm at 10 seconds intervals for five minutes in a THERMOMax plate reader at ambient temperature. When a lower concentration of factor Xa was needed in the factor Xa assay, the enzyme kinetics were measured for fifteen minutes (0.2 nM factor Xa) or thirty minutes (0.1 nM factor Xa) at ambient temperature.

15

Calculation of K_i of the Test compounds:

Enzyme initial rates were calculated as mOD/min based on the first two minutes readings. The IC_{50} values were determined by fitting the data to the log-logit equation (linear) or the Morrison equation (non-linear) with an EXCEL spread-sheet. K_i values were then obtained by dividing the IC_{50} by 2. Routinely, K_i (factor Xa) values less than 3 nM were calculated from the Morrison equation.

Compounds of the invention, when tested in this assay, demonstrated the selective ability to inhibit human factor Xa and human thrombin.

25

EXAMPLE 9

(In vitro assay for Human Prothrombinase)

This assay demonstrates the ability of the compounds of the invention to inhibit prothrombinase. Prothrombinase (PTase) catalyzes the activation of prothrombin to yield fragment 1.2 plus thrombin with meizothrombin as the intermediate. This assay is an end point assay. Activity of the prothrombinase is measured by activity of thrombin (one of the reaction products) or by the amount

30

of thrombin formed/time based on a thrombin standard curve (nM vs mOD/min).
For determination of IC₅₀ (PTase) of the compounds of the invention, PTase activity was expressed by thrombin activity (mOD/min).

Materials:

5 Enzymes:

1. Human factor Va (Haematologic Technologies Inc., Cat# HCVA-0110) working solution: 1.0 mg/mL in 50% glycerol, 2 mM CaCl₂, stored at -20°C.
2. Human factor Xa (Enzyme Res. Lab. cat# HFXa1011) working solution:
10 0.281 mg/mL in assay buffer (without BSA), stored at -80°C.
3. Human prothrombin (FII) (Enzyme Res. Lab., Cat# HP1002) working solution:
Diluted FII to 4.85 mg/mL in assay buffer (without BSA), stored at -80°C.

Phospholipid (PCPS) vesicles:

- 15 PCPS vesicles (80%PC, 20%PS) were prepared by modification of the method reported by Barenholz *et al.*, *Biochemistry* (1977), Vol. 16, pp. 2806-2810.

Phosphatidyl serine (Avanti Polar Lipids, Inc., Cat#840032):

- 10 10 mg/mL in chloroform, purified from brain, stored -20°C under nitrogen or argon.

Phosphatidyl Choline (Avanti Polar Lipids, Inc., Cat# 850457):

50 mg/ml in chloroform, synthetic 16:0-18:1 Palmitoyl-Oleoyl, stored at -20°C under nitrogen or argon.

Spectrozyme-TH (American Diagnostica Inc., Cat# 238L, 50 µmoles, stored at

- 25 ambient temperature) working solution: Dissolved 50 µmoles in 10 mL dH₂O.

BSA (Sigma Chem Co., Cat# A-7888, FractionV, RIA grade).

Assay buffer: 50 mM TrisHCl, pH 7.5, 150 mM NaCl, 2.5 mM CaCl₂, 0.1% PEG 6000

- 30 (BDH), 0.05% BSA (Sigma, Fr.V, RIA grade).

For one plate assay, prepare the following working solutions:

1. Prothrombinase complex:

- (a) 100 μ M PCPS (27.5 μ L of PCPS stock (4.36 mM) diluted to final 1200 μ L with assay buffer.
- (b) 25 nM Human factor Va: 5.08 μ L of Va stock (1 mg/mL) was diluted to final 1200 μ L with assay buffer.
- 5 (c) 5 pM Human factor Xa: Dilute factor Xa stock (0.281 mg/mL) 1:1,220,000 with assay buffer. Prepare at least 1200 μ L.

Combine equal volumes (1100 μ L) of each component in the order of PCPS, Va and Xa. Use immediately or store in ice (bring to ambient temperature before use).

- 10 2. 6 μ M Human prothrombin (FII): dilute 124 μ L of FII stock (4.85 mg/mL) to final 1400 μ L with assay buffer.
3. 20 mM EDTA/Assay buffer: 0.8 mL of 0.5 M EDTA (pH 8.5) plus 19.2 mL assay buffer.
4. 0.2 mM Spectrozyme-TH/EDTA buffer: 0.44 mL of SPTH stock (5 mM) plus 10.56 mL of 20 mM EDTA/assay buffer.
- 15 5. Test compounds (compounds of the invention):
Prepare a working solution (5X) from 10 mM stock (DMSO) and make a series of 1:3 dilution. Compounds were assayed at 6 concentrations in duplicate.

20

Assay conditions and procedure:

Prothrombinase reaction was performed in final 50 μ L of mixture containing PTase (20 μ M PCPS, 5 nM hFVa, and 1 pM hFXa), 1.2 μ M human factor II and
25 varied concentration of the test compounds (5 μ M to 0.021 μ M or lower concentration range). Reaction was started by addition of PTase and incubated for 6 minutes at ambient temperature. Reaction was stopped by addition of EDTA/buffer to final 10 mM. Activity of thrombin (product) was then measured in the presence of 0.1 mM of Spectrozyme-TH as substrate at 405 nm for 5
30 minutes (10 seconds intervals) at ambient temperature in a THEROmax microplate reader. Reactions were performed in 96-well microtiter plates.

In the first step of the assay, 10 μ L of diluted test compound (5X) or buffer was added to the plates in duplicate. Then 10 μ L of prothrombin (hFII) (5X) was added to each well. Next 30 μ L PTase was added to each well, mix for about 30 seconds. The plates were then incubated at ambient temperature for 6 minutes.

5

In the next step, 50 μ L of 20 mM EDTA (in assay buffer) was added to each well to stop the reaction. The resulting solutions were then mixed for about 10 seconds. Then 100 μ L of 0.2 mM spectrozyme was added to each well. The thrombin reaction rate was then measured at 405 nm for 5 minutes at 10 seconds intervals in a Molecular Devices microplate reader.

10

Calculations:

Thrombin reaction rate was expressed as mOD/min. using OD readings from the five minute reaction. IC₅₀ values were calculated with the log-logit curve fit program.

15

The compounds of the invention demonstrated the ability to inhibit pro-thrombinase when tested in this assay.

20

EXAMPLE 10

(In vivo assay)

The following assay demonstrates the ability of the compounds to act as anti-coagulants.

25

Male rats (250-330 g) were anesthetized with sodium pentobarbital (90 mg/kg, i.p.) and prepared for surgery. The left carotid artery was cannulated for the measurement of blood pressure as well as for taking blood samples to monitor clotting variables (prothrombin time (PT) and activated partial thromboplastin time (aPTT)). The tail vein was cannulated for the purpose of administering the test compounds (*i.e.*, the compounds of the invention and

30

standards) and the thromboplastin infusion. The abdomen was opened *via* a mid-line incision and the abdominal vena cava was isolated for 2-3 cm distal to the renal vein. All venous branches in this 2-3 cm segment of the abdominal vena cava were ligated. Following all surgery, the animals were allowed to
5 stabilize prior to beginning the experiment. Test compounds were administered as an intravenous bolus ($t=0$). Three minutes later ($t=3$), a 5-minute infusion of thromboplastin was begun. Two minutes into the infusion ($t=5$), the abdominal vena cava was ligated at both the proximal and distal ends. The vessel was left in place for 60 minutes, after which it was excised from the animal, slit open, the
10 clot (if any) carefully removed, and weighed. Statistical analysis on the results was performed using a Wilcoxin-matched-pairs signed rank test.








The compounds of the invention, when tested in this assay, demonstrated the ability to
15 inhibit the clotting of the blood.

* * * * *

While the present invention has been described with reference to the specific
20 embodiments thereof, it should be understood by those skilled in the art that various changes may be made and equivalents may be substituted without departing from the true spirit and scope of the invention. In addition, many modifications may be made to adapt a particular situation, material, composition of matter, process, process step or steps, to the objective, spirit and scope of the
25 present invention. All such modifications are intended to be within the scope of the claims appended hereto.

WHAT IS CLAIMED IS:

1. A compound selected from the group consisting of the following formulae:



5

wherein:

A is $-C(R^8)=$ or $-N=$ (where R^8 is hydrogen, alkyl or halo);

Z^1 is $-O-$, $-N(R^9)-$, $-CH_2O-$ or $-S(O)_n-$ (where n is 0 to 2);

Z^2 is $-O-$, $-N(R^9)-$, $-OCH_2-$ or $-S(O)_n-$ (where n is 0 to 2);

- 10 R^1 and R^4 are each independently hydrogen, halo, alkyl, nitro, $-OR^9$, $-C(O)OR^9$, $-C(O)N(R^9)R^{10}$, $-N(R^9)R^{10}$, $-N(R^9)C(O)R^9$, or $-N(H)S(O)_2R^{11}$;

-53-

R^2 is $-C(NH)NH_2$, $-C(NH)N(H)OR^9$, $-C(NH)N(H)C(O)OR^{11}$, $-C(NH)N(H)C(O)R^9$, $-C(NH)N(H)S(O)_2R^{11}$, or $-C(NH)N(H)C(O)N(H)R^9$;

R^3 is hydrogen, halo, alkyl, haloalkyl, nitro, ureido, guanidino, $-OR^9$, $-C(NH)NH_2$, $-C(NH)N(H)OR^9$, $-C(O)N(R^9)R^{10}$, $-R^{12}-C(O)N(R^9)R^{10}$, $-CH(OH)C(O)N(R^9)R^{10}$, $-N(R^9)R^{10}$, $-R^{12}-N(R^9)R^{10}$, $-C(O)OR^9$, $-R^{12}-C(O)OR^9$, $-N(R^9)C(O)R^9$, (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl), (1,2)-imidazolyl (optionally substituted by alkyl), or (1,2)-imidazolinyl (optionally substituted by alkyl);

10 R⁵ and R⁶ are independently hydrogen, halo, alkyl, haloalkyl, nitro, -N(R⁹)R¹⁰, -C(O)OR⁹, -C(O)N(R⁹)R¹⁰, -C(O)N(R⁹)CH₂C(O)N(R⁹)R¹⁰, -N(R⁹)C(O)N(R⁹)R¹⁰, -N(R⁹)C(O)R¹⁰, -N(R⁹)S(O)₂R¹¹, or -N(R⁹)C(O)N(R⁹)-CH₂C(O)N(R⁹)R¹⁰;

R^7 is $-X-CH_2-[C(R^{13})H]_p-C(R^{13})H_2$ or $-X-C([C(R^{13})H]_p-C(R^{13})H_2)_2H$ where:

p is 0 to 5;

15 X is -O-, -S(O)_n- (where n is 0 to 2), or -N(R⁹)-;

and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -[CH₂]_n-OC(O)R⁹, -[CH₂]_n-SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or -[CH₂]_n-N(R⁹)C(O)N(R⁹)R¹⁰ (where each n is independently 0 to 2);

20 $N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

or R⁷ is selected from the group consisting of the following formulae:



where

q is 0 to 4;

25 p is 0 to 5;

each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -(CH₂)_n-OC(O)R⁹, -[CH₂]_n-SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or

-54-

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0 to 2);

and

each R^{14} is $-\text{X-[CH}_2\text{]}_r-$ or $-\text{X-CH}_2\text{-[C(R}^{13}\text{)H]}_r-$, where:

each r is independently 0 to 5, and

each X is $-\text{O}-$, $-\text{S(O)}_n-$ (where n is 0 to 2), or $-\text{N(R}^9\text{)}-$; and

each R^{13} is independently $-\text{[CH(OR}^9\text{)]}_m\text{-CH}_2\text{-OR}^9$ (where m is 1 to 4),

$-\text{[CH}_2\text{]}_n\text{-OR}^9$, $-\text{[CH}_2\text{]}_n\text{-SR}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)R}^{10}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)R}^9$,

$-\text{[CH}_2\text{]}_n\text{-SC(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)R}^9$, $-\text{[CH}_2\text{]}_n\text{-OC(O)OR}^{11}$,

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)OR}^{11}$, $-\text{[CH}_2\text{]}_n\text{-OC(O)N(R}^9\text{)R}^{10}$, or

$-\text{[CH}_2\text{]}_n\text{-N(R}^9\text{)C(O)N(R}^9\text{)R}^{10}$ (where each n is independently 0 to 2);


each R^9 and R^{10} is independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl);

each R^{11} is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl); and

each R^{12} is independently an alkylene or alkylidene chain;

as a single stereoisomer or a mixture thereof; or a pharmaceutically acceptable salt thereof.

2. The compound of Claim 1 selected from formula (I):



as a single stereoisomer or a mixture thereof; or a pharmaceutically acceptable salt thereof.

3. The compound of Claim 2 wherein:

A is -N=;

Z¹ is -O-, -CH₂O- or -S(O)_n- (where n is 1);

10 Z² is -O-, -OCH₂- or -S(O)_n- (where n is 1);

R¹ and R⁴ are each independently hydrogen, halo, alkyl or -OR⁹;

R² is -C(NH)NH₂, -C(NH)N(H)S(O)₂R¹¹ or -C(NH)N(H)C(O)N(H)R⁹;

R³ is ureido, guanidino, -N(R⁹)R¹⁰, -N(R⁹)C(O)R⁹, (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl), (1,2)-imidazolyl (optionally substituted by

15 alkyl) or (1,2)-imidazolyl (optionally substituted by alkyl);

R⁵ and R⁶ are independently hydrogen, halo, alkyl or haloalkyl;

R⁷ is -X-CH₂-[C(R¹³)H]_p-C(R¹³)H₂ where:

p is 0 to 5;

X is -O-, -S(O)_n- (where n is 0 to 2), or -N(R⁹)-;

20 and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -[CH₂]_n-OC(O)R⁹, -[CH₂]_n-SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or -[CH₂]_n-N(R⁹)C(O)N(R⁹)R¹⁰ (where each n is independently 0 to 2);

25 each R⁹ and R¹⁰ is independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl,

-56-

hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl); and

each R¹¹ is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

4. The compound of Claim 3 wherein:

A is -N=;

Z¹ is -O-;

15 Z² is -O-;

R¹ is hydrogen or -OR⁹;

R² is -C(NH)NH₂;

R³ is (1,2)-tetrahydropyrimidinyl (optionally substituted by methyl), (1,2)-imidazolyl (optionally substituted by methyl) or (1,2)-imidazoliny (optionally substituted by methyl);

R⁴ is hydrogen;

R⁵ and R⁶ are each halo;

R⁷ is -X-CH₂-[C(R¹³)H]_p-C(R¹³)H₂ where:

p is 0 to 5;

25 X is -O-;

and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹, -[CH₂]_n-N(R⁹)R¹⁰, -[CH₂]_n-OC(O)R⁹, -[CH₂]_n-SC(O)R⁹, -[CH₂]_n-N(R⁹)C(O)R⁹, -[CH₂]_n-OC(O)OR¹¹, -[CH₂]_n-N(R⁹)C(O)OR¹¹, -[CH₂]_n-OC(O)N(R⁹)R¹⁰, or -[CH₂]_n-N(R⁹)C(O)N(R⁹)R¹⁰ (where each n is independently 0 to 2);

each R⁹ and R¹⁰ is independently hydrogen or alkyl; and

each R¹¹ is alkyl or aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy,

-57-

alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

5. The compound of Claim 4 wherein:

5 A is -N=;

Z¹ is -O-;

Z² is -O-;

R¹ is -OR⁹;

R² is -C(NH)NH₂;

10 R³ is (1,2)-imidazolyl (optionally substituted by methyl) or (1,2)-imidazoliny
(optionally substituted by methyl);

R⁴ is hydrogen;

R⁵ and R⁶ are both fluoro;

R⁷ is -X-CH₂-[C(R¹³)H]_p-C(R¹³)H₂ where:

15 p is 0 to 5;

X is -O-;

and each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -
[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹ or -[CH₂]_n-N(R⁹)R¹⁰ (where each n is
independently 0 to 2); and

20 each R⁹ and R¹⁰ is independently hydrogen or alkyl.

6. The compound of Claim 5 wherein:

R⁷ is -O-CH₂-C(OH)H-C(OH)H₂.

25 7. The compound of Claim 6, namely, 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-(2,3-dihydroxypropoxy)pyridin-2-yl]oxy]-4-hydroxybenzamidine.

8. The compound of Claim 2 wherein:

30 A is -N=;

Z¹ is -O-, -CH₂O- or -S(O)_n- (where n is 0 to 2);

Z² is -O-, -OCH₂- or -S(O)_n- (where n is 0 to 2);

R¹ and R⁴ are each independently hydrogen, halo or -OR⁹;

-58-

R^2 is $-C(NH)NH_2$, $-C(NH)N(H)S(O)_2R^{11}$ or $-C(NH)N(H)C(O)N(H)R^9$;

R^3 is ureido, guanidino, $-N(R^9)R^{10}$, $-N(R^9)C(O)R^9$, (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl), (1,2)-imidazolyl (optionally substituted by alkyl), or (1,2)-imidazolinyl (optionally substituted by alkyl);

5 R^5 and R^6 are independently hydrogen, halo, alkyl or haloalkyl;

R^7 is selected from the group consisting of the following formulae:



10 where

q is 0 to 4;

p is 0 to 5;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -

$[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-(CH_2)_n-OC(O)R^9$,

15 $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$,

$-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or

$-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

and

each R^{14} is $-X-[CH_2]_r$ or $-X-CH_2-[C(R^{13})H]_r$, where:

20 each r is independently 0 to 5, and

each X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4),

$-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-[CH_2]_n-OC(O)R^9$,

$-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$,

25 $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or

$-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

each R^9 and R^{10} is independently hydrogen, alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or

30

-59-


dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl);

- 5 each R^{11} is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl, hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
- 10 dialkylaminocarbonyl).


9. The compound of Claim 8 wherein:

- A is $-N=$;
- 15 Z^1 is $-O-$;
- Z^2 is $-O-$;
- R^1 is hydrogen or $-OR^9$;
- R^2 is $-C(NH)NH_2$;
- R^3 is (1,2)-tetrahydropyrimidinyl (optionally substituted by methyl), (1,2)-imidazolyl
- 20 (optionally substituted by methyl), or (1,2)-imidazoliny (optionally substituted by methyl);
- R^4 is hydrogen;
- R^5 and R^6 are each halo;
- R^7 is selected from the group consisting of the following formulae:

25



and



where

q is 0 to 4;

p is 0 to 5;

-60-

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-(CH_2)_n-OC(O)R^9$, $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$, $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or $-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);
and

each R^{14} is $-X-[CH_2]_r$ or $-X-CH_2-[C(R^{13})H]_r$, where:

each r is independently 0 to 5, and

each X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-[CH_2]_n-OC(O)R^9$, $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$, $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or $-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

each R^9 and R^{10} is independently hydrogen or alkyl; and

each R^{11} is independently alkyl or aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy, alkoxy carbonyl, aminocarbonyl, monoalkylaminocarbonyl, or dialkylaminocarbonyl).

10. The compound of Claim 9 wherein:

A is $-N=$;

Z^1 is $-O-$;

25 Z^2 is $-O-$;

R^1 is $-OR^9$;

R^2 is $-C(NH)NH_2$;


R^3 is (1,2)-imidazolyl (optionally substituted by methyl), or (1,2)-imidazolyl (optionally substituted by methyl);

30 R^4 is hydrogen;

R^5 and R^6 are both fluoro;

R^7 is selected from the group consisting of the following formulae:

-61-



5

where

q is 0 to 4;

p is 0 to 5;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), - $[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is

10

independently 0 to 2); and


each R^{14} is $-X-[CH_2]_r-$ or $-X-CH_2-[C(R^{13})H]_r-$, where:

each r is independently 0 to 5, and

each X is independently $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$;

and

15

each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$; andeach R^9 and R^{10} is independently hydrogen or alkyl.11. The compound of Claim 10 wherein R^7 is:

20

where

q is 0 to 4;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$ or $-[CH_2]_n-N(R^9)R^{10}$ (where each n is independently 0 to 2); and

25

each R^{14} is $-X-[CH_2]_r-$ or $-X-CH_2-[C(R^{13})H]_r-$, where:


each r is independently 0 to 5, and

-62-

each X is independently -O-, -S(O)_n- (where n is 0 to 2), or -N(R⁹)-; and
 each R¹³ is independently -[CH(OR⁹)]_m-CH₂-OR⁹ (where m is 1 to 4),
 -[CH₂]_n-OR⁹, -[CH₂]_n-SR⁹ or -[CH₂]_n-N(R⁹)R¹⁰; and
 each R⁹ and R¹⁰ is independently hydrogen or alkyl.

5

12. The compound of Claim 11 wherein R⁷ is:



where

10 q is 0 to 4;


each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹,
 -[CH₂]_n-SR⁹ or -[CH₂]_n-N(R⁹)R¹⁰ (where each n is independently 0 to 2); and
 each R¹⁴ is -X-[CH₂]_r- where:

r is 0 to 5, and

15 X is -O-, -S(O)_n- (where n is 0 to 2), or -N(R⁹)-; and
 each R⁹ and R¹⁰ is independently hydrogen or alkyl.

13. The compound of Claim 12 wherein R⁷ is:

20



where

q is 0 to 4;

25 each R¹³ is independently -[C(OR⁹)H]_m-CH₂-OR⁹ (where m is 1 to 4), -[CH₂]_n-OR⁹
 or -[CH₂]_n-N(R⁹)R¹⁰ (where each n is independently 0 to 2); and

-63-

each R^{14} is $-X-[CH_2]_r$ where:


r is 0 or 1, and

X is $-O-$; and

each R^9 and R^{10} is independently hydrogen or alkyl.

5

14. The compound of Claim 13 wherein R^7 is:










where R^{14} is $-O-$; q is 3 and one R^{13} is in the 4-position of the tetrahydrofuranyl ring and is hydroxy, the second R^{13} is in the 2-position of the tetrahydrofuranyl ring and
 10 is 2,3-dihydroxyethyl and the third R^{13} is in the 5-position of the tetrahydrofuranyl ring and is ethoxy.

15. The compound of Claim 14, namely, 3-[[3,5-difluoro-6-(3-(1-methylimidazolin-2-yl)phenoxy)-4-[[2-(1,2-dihydroxyethyl)-4-hydroxy-5-ethoxytetrahydrofuran-3-yl]oxy]pyridin-2-yl]oxy]-4-hydroxybenzamidine.

16. A pharmaceutical composition useful in treating a human having a disease-state characterized by thrombotic activity, which composition comprises a therapeutically effective amount of a compound selected from the group
 20 consisting of the following formulae:

-64-



wherein:

- 5 A is $-\text{C}(\text{R}^8)=$ or $-\text{N}=$ (where R^8 is hydrogen, alkyl or halo);
 Z^1 is $-\text{O}-$, $-\text{N}(\text{R}^9)-$, $-\text{CH}_2\text{O}-$ or $-\text{S}(\text{O})_n-$ (where n is 0 to 2);
 Z^2 is $-\text{O}-$, $-\text{N}(\text{R}^9)-$, $-\text{OCH}_2-$ or $-\text{S}(\text{O})_n-$ (where n is 0 to 2);
 R^1 and R^4 are each independently hydrogen, halo, alkyl, nitro, $-\text{OR}^9$, $-\text{C}(\text{O})\text{OR}^9$,
 $-\text{C}(\text{O})\text{N}(\text{R}^9)\text{R}^{10}$, $-\text{N}(\text{R}^9)\text{R}^{10}$, $-\text{N}(\text{R}^9)\text{C}(\text{O})\text{R}^9$, or $-\text{N}(\text{H})\text{S}(\text{O})_2\text{R}^{11}$;
- 10 R^2 is $-\text{C}(\text{NH})\text{NH}_2$, $-\text{C}(\text{NH})\text{N}(\text{H})\text{OR}^9$, $-\text{C}(\text{NH})\text{N}(\text{H})\text{C}(\text{O})\text{OR}^{11}$, $-\text{C}(\text{NH})\text{N}(\text{H})\text{C}(\text{O})\text{R}^9$,
 $-\text{C}(\text{NH})\text{N}(\text{H})\text{S}(\text{O})_2\text{R}^{11}$, or $-\text{C}(\text{NH})\text{N}(\text{H})\text{C}(\text{O})\text{N}(\text{H})\text{R}^9$;

-65-

R^3 is hydrogen, halo, alkyl, haloalkyl, nitro, ureido, guanidino, $-OR^9$, $-C(NH)NH_2$,
 $-C(NH)N(H)OR^9$, $-C(O)N(R^9)R^{10}$, $-R^{12}-C(O)N(R^9)R^{10}$, $-CH(OH)C(O)N(R^9)R^{10}$,
 $-N(R^9)R^{10}$, $-R^{12}-N(R^9)R^{10}$, $-C(O)OR^9$, $-R^{12}-C(O)OR^9$, $-N(R^9)C(O)R^9$,
 (1,2)-tetrahydropyrimidinyl (optionally substituted by alkyl), (1,2)-imidazolyl
 5 (optionally substituted by alkyl), or (1,2)-imidazolinyl (optionally substituted
 by alkyl);

R^5 and R^6 are independently hydrogen, halo, alkyl, haloalkyl, nitro, $-N(R^9)R^{10}$,
 $-C(O)OR^9$, $-C(O)N(R^9)R^{10}$, $-C(O)N(R^9)CH_2C(O)N(R^9)R^{10}$,
 $-N(R^9)C(O)N(R^9)R^{10}$, $-N(R^9)C(O)R^{10}$, $-N(R^9)S(O)_2R^{11}$, or -
 10 $N(R^9)C(O)N(R^9)-CH_2C(O)N(R^9)R^{10}$;

R^7 is $-X-CH_2-[C(R^{13})H]_p-C(R^{13})H_2$ or $-X-C([C(R^{13})H]_p-C(R^{13})H_2)_2H$ where:

p is 0 to 5;

X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$;

and each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -

15 $[CH_2]_n-OR^9$, $[CH_2]_n-SR^9$, $[CH_2]_n-N(R^9)R^{10}$, $[CH_2]_n-OC(O)R^9$, $[CH_2]_n-$
 $SC(O)R^9$, $[CH_2]_n-N(R^9)C(O)R^9$, $[CH_2]_n-OC(O)OR^{11}$, $[CH_2]_n-$
 $N(R^9)C(O)OR^{11}$, $[CH_2]_n-OC(O)N(R^9)R^{10}$, or $[CH_2]_n-$
 $N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);

or R^7 is selected from the group consisting of the following formulae:



20

where

q is 0 to 4;

p is 0 to 5;

each R^{13} is independently $-[C(OR^9)H]_m-CH_2-OR^9$ (where m is 1 to 4), -

25 $[CH_2]_n-OR^9$, $[CH_2]_n-SR^9$, $[CH_2]_n-N(R^9)R^{10}$, $[CH_2]_n-OC(O)R^9$,
 $[CH_2]_n-SC(O)R^9$, $[CH_2]_n-N(R^9)C(O)R^9$, $[CH_2]_n-OC(O)OR^{11}$,
 $[CH_2]_n-N(R^9)C(O)OR^{11}$, $[CH_2]_n-OC(O)N(R^9)R^{10}$, or
 $[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0 to 2);
 and

-66-

each R^{14} is $-X-[CH_2]_r-$ or $-X-CH_2-[C(R^{13})H]_r-$, where:

each r is independently 0 to 5, and

each X is $-O-$, $-S(O)_n-$ (where n is 0 to 2), or $-N(R^9)-$; and

each R^{13} is independently $-[CH(OR^9)]_m-CH_2-OR^9$ (where m is 1 to 4),

5 $-[CH_2]_n-OR^9$, $-[CH_2]_n-SR^9$, $-[CH_2]_n-N(R^9)R^{10}$, $-[CH_2]_n-OC(O)R^9$,
 $-[CH_2]_n-SC(O)R^9$, $-[CH_2]_n-N(R^9)C(O)R^9$, $-[CH_2]_n-OC(O)OR^{11}$,
 $-[CH_2]_n-N(R^9)C(O)OR^{11}$, $-[CH_2]_n-OC(O)N(R^9)R^{10}$, or
 $-[CH_2]_n-N(R^9)C(O)N(R^9)R^{10}$ (where each n is independently 0
to 2);

10 each R^9 and R^{10} is independently hydrogen, alkyl, aryl (optionally substituted by
halo, alkyl, hydroxy, alkoxy, aralkoxy, amino, dialkylamino, monoalkylamino,
nitro, carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl,
hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro,
15 carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
dialkylaminocarbonyl);

each R^{11} is alkyl, aryl (optionally substituted by halo, alkyl, hydroxy, alkoxy,
aralkoxy, amino, dialkylamino, monoalkylamino, nitro, carboxy,
alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
20 dialkylaminocarbonyl), or aralkyl (optionally substituted by halo, alkyl, aryl,
hydroxy, alkoxy, aralkyl, amino, dialkylamino, monoalkylamino, nitro,
carboxy, alkoxycarbonyl, aminocarbonyl, monoalkylaminocarbonyl, or
dialkylaminocarbonyl); and

each R^{12} is independently an alkylene or alkylidene chain;

25 as a single stereoisomer or a mixture thereof; or a pharmaceutically acceptable salt
thereof; and a pharmaceutically acceptable excipient.

17. Use of one or several compounds of claim 1 for the manufacture of a
medicament for the treatment of a disease characterized by thrombotic activity.

INTERNATIONAL SEARCH REPORT

International Application No
PCT/IB 99/02067

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C07D401/12 C07D405/14 A61K31/44 A61P7/02

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C07D A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 98 11094 A (SCHERING AG) 19 March 1998 (1998-03-19) claim 1; examples ---	1-17
X	WO 98 15547 A (SCHERING AG) 16 April 1998 (1998-04-16) claim 1; examples ---	1-17
X	WO 96 28427 A (BERLEX LAB ;BUCKMAN BRAD O (US); DAVEY DAVID D (US); GUILFORD WILL) 19 September 1996 (1996-09-19) claim 1; examples ---	1-17
X	WO 97 29067 A (BERLEX LAB) 14 August 1997 (1997-08-14) cited in the application claim 1; examples ---	1-17
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

19 April 2000

Date of mailing of the international search report

09/05/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

De Jong, B

INTERNATIONAL SEARCH REPORT

International Application No

PCT/IB 99/02067

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	<p>PHILLIPS, GARY B. ET AL: "Discovery of N-'2-'5-'Amino(imino)methyl-2-hydroxyphenoxy-3,5- difluoro-6-'3-(4,5-dihydro-1-methyl-1H-imidazol-2-yl)phenoxy!pyridin-4-yl!-N-methylglycine (ZK-807834): A Potent, Selective, and Orally Active Inhibitor of the Blood Coagulation Enzyme Factor Xa1"</p> <p>J. MED. CHEM. (1998), 41(19), 3557-3562 , XP002136093</p> <p>-----</p>	1-17

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/IB 99/02067

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9811094 A	19-03-1998	US 6008234 A	28-12-1999
		AU 4384397 A	02-04-1998
		CN 1234798 A	10-11-1999
		CZ 9900861 A	11-08-1999
		EP 0929547 A	21-07-1999
		NO 991206 A	11-05-1999
		PL 332092 A	30-08-1999
		SK 29599 A	06-08-1999
WO 9815547 A	16-04-1998	US 6004985 A	21-12-1999
		AU 4624097 A	05-05-1998
		CN 1233247 A	27-10-1999
		CZ 9901227 A	15-09-1999
		EP 0934310 A	11-08-1999
		NO 991594 A	08-06-1999
		PL 332602 A	27-09-1999
		US 6034084 A	07-03-2000
WO 9628427 A	19-09-1996	US 5691364 A	25-11-1997
		AU 707323 B	08-07-1999
		AU 5299496 A	02-10-1996
		CA 2214685 A	19-09-1996
		EP 0813525 A	29-12-1997
		US 6034103 A	07-03-2000
		US 5883100 A	16-03-1999
		US 5877181 A	02-03-1999
WO 9729067 A	14-08-1997	US 5889005 A	30-03-1999
		US 5994375 A	30-11-1999
		AU 715992 B	17-02-2000
		AU 1454897 A	28-08-1997
		CA 2245925 A	14-08-1997
		CN 1211232 A	17-03-1999
		CZ 9802550 A	13-01-1999
		EP 0888271 A	07-01-1999
		HU 9902017 A	28-09-1999
		NO 983671 A	11-09-1998
		PL 328412 A	18-01-1999
		SK 109298 A	11-02-1999